

How can I find the closest value to a given number in a vector?

Authored by
stats writer

June 26, 2024

RECOMMENDED CITATION

stats writer (2024). *How can I find the closest value to a given number in a vector?*. PSYCHOLOGICAL SCALES. Retrieved from <https://scales.arabpsychology.com/?p=154051>

The process of finding the closest value to a given number in a vector involves searching through the vector to determine the element that is closest in value to the specified number. This can be achieved by calculating the absolute difference between each element in the vector and the given number, and then selecting the element with the smallest difference. This method allows for efficient and accurate retrieval of the closest value in a vector.

R: Find Closest Value in Vector

You can use the following basic syntax to find the closest value between elements of two vectors in R:

```
#define cut points
```

```
cuts <- c(-Inf, vector2-diff(vector2)/2, Inf)
```

```
#for each value in vector1, find closest value in vector2
```

```
cut(vector1, breaks=cuts, labels=vector2)
```

The following example shows how to use this syntax in practice.

Example: Find Closest Value in Vector in R

Suppose we have the following two vectors in R:

```
#define vectors
```

```
vector1 <- c(1, 2, 3, 4, 5, 6, 7, 8, 9, 10)
```

```
vector2 <- c(3, 5, 8, 11)
```

Now suppose that for each value in the first vector, we would like to find the closest value in the second vector.

We can use the following syntax to do so:

```
#define cut points
```

```
cuts <- c(-Inf, vector2-diff(vector2)/2, Inf)
```

```
#for each value in vector1, find closest value in vector2  
cut(vector1, breaks=cuts, labels=vector2)
```

```
3 3 3 3 5 5 8 8 8 11
```

Here's how to interpret the output:

For the first value in vector1 (1), the closest value in vector2 is 3. For the second value in vector1 (2), the closest value in vector2 is 3. For the third value in vector1 (3), the closest value in vector2 is 3. For the fourth value in vector1 (4), the closest value in vector2 is 3. For the fifth value in vector1 (5), the closest value in vector2 is 5.

And so on.

Note: This method assumes that the values in the second vector are strictly increasing. If they aren't already, you may need to first sort the second vector.

ARABPSYCHOLOGY.COM