

# How to Calculate and Round the Average in Google Sheets

Authored by  
**stats writer**

January 25, 2026

## RECOMMENDED CITATION

stats writer (2026). *How to Calculate and Round the Average in Google Sheets*.

PSYCHOLOGICAL SCALES. Retrieved from <https://scales.arabpsychology.com/?p=127582>

Google Sheets is a cornerstone tool for data professionals and analysts, enabling sophisticated calculations and detailed data manipulation across various datasets. A particularly common requirement is the ability to calculate the average of a collection of numbers and simultaneously ensure the result is presented cleanly, often by using rounding. This guide focuses on combining the power of the AVERAGE function and the ROUND function to achieve precise and whole numerical outputs.

The standard process involves first aggregating the specified range of values using the AVERAGE function. The result, which is often a floating-point number with many decimal places, is then passed as an argument into the ROUND function. This nesting technique allows for immediate calculation and formatting in a single, efficient cell entry. Mastering this combination is essential for generating reports that demand simplified averages, ensuring that data analysis and subsequent decision-making processes are both quick and grounded in accurate, easily digestible figures.

## Google Sheets: Calculate Average with Rounding

### The Power of Data Aggregation in Google Sheets

Analyzing large datasets often requires summarizing information into key statistics. The average, or arithmetic mean, is arguably the most fundamental statistical measure used in reporting, giving stakeholders a centralized value around which the data points cluster. While Google Sheets makes calculating the average effortless using the built-in AVERAGE function, the raw output frequently includes a long string of decimal places that can clutter reports and obscure the true meaning of the data.

The goal, therefore, is not merely to calculate the mean but to present it in a format that aids comprehension and maintains computational integrity. For many business or academic applications, presenting an average rounded to a specific number of decimal places or, crucially, to the nearest whole integer, is a requirement. This necessity for clean, precise reporting drives the need to integrate rounding functionality directly into the averaging calculation.

### Core Functions Explained: AVERAGE and ROUND

To successfully combine these operations, one must understand the syntax and purpose of the two primary functions involved. The first, the AVERAGE function, takes a range of cells or a list of numbers as its input and returns the mathematical average of those numerical values. Its basic structure is straightforward: `=AVERAGE(value1, )`, where `value1` is typically a cell range like **A2:A14**.

The second essential component is the ROUND function. This function is designed specifically for

rounding a number to a specified number of places. It requires two arguments: the number to be rounded (which will be the output of our AVERAGE calculation) and the number of places to round to. The syntax is `=ROUND(number, places)`. By nesting the AVERAGE function inside the ROUND function, we create a powerful formula that performs two operations simultaneously, ensuring the calculation occurs before the formatting.

## Implementing the Combined Formula Structure

There are generally two widely accepted methods for calculating the average value of a data range in Google Sheets and simultaneously formatting the result using rounding. Both methods rely on the principle of function nesting, where the result of the inner function (AVERAGE) feeds directly into the outer function (ROUND).

The choice between the methods depends entirely on the required level of precision for the final output: whether the report requires exact values rounded to a few decimal places, or if it mandates a simple, non-fractional whole number.

### Method 1: Rounding the Average to Specific Decimal Places

If your reporting requirements necessitate maintaining some level of fractional precision--for instance, showing financial results to two decimal places for currency, or scientific measurements to three significant digits--you should use the ROUND function with a specified positive integer for the second argument.

This method ensures that the calculated mean value remains highly accurate while being easily readable. The second parameter in the ROUND function determines exactly how many digits will appear after the decimal point in the final averaged output.

The structure for this approach is as follows:

```
=ROUND(AVERAGE(A2:A14), 3)
```

This precise formula calculates the average value of the cells in the range **A2:A14**. Subsequently, the function takes this raw average and rounds the result specifically to **3 decimal places**, balancing accuracy with presentation requirements.

### Detailed Example 1: Controlling Precision

To illustrate Method 1, consider a column of data representing daily sales figures or measurement readings. Before applying the formula, it is helpful to visualize the dataset used in this calculation. We will use the following collection of values in Google Sheets, which, when averaged, results in

approximately **20.0769**:

C2    ▾    *fx* =AVERAGE(A2:A14)

	A	B	C	D
1	<b>Values</b>		<b>Average</b>	
2	14		20.07692308	
3	19			
4	30			
5	24			
6	25			
7	25			
8	23			
9	18			
10	15			
11	13			
12	11			
13	10			
14	34			
15				
16				
17				

We can input the combined formula into an empty cell, such as **D2**, to calculate the average of the data range **A2:A14** and ensure it is rounded to three decimal places. This action is critical for reports that must adhere to specific formatting standards, such as those used in engineering or finance where three digits of precision are often standard practice.

The exact formula entered into cell D2 is:

**=ROUND(AVERAGE(A2:A14), 3)**

The immediate visual output of this formula execution is demonstrated in the subsequent screenshot, confirming the successful calculation and rounding operation:

D2     $\nabla$  |  $fx$  =ROUND(AVERAGE(A2:A14), 3)

	A	B	C	D
1	<b>Values</b>		<b>Average</b>	<b>Average Rounded to 3 Decimals</b>
2	14		20.07692308	20.077
3	19			
4	30			
5	24			
6	25			
7	25			
8	23			
9	18			
10	15			
11	13			
12	11			
13	10			
14	34			
15				
16				

Upon execution, the formula returns the average value of the cells within the specified range **A2:A14**, accurately rounded to 3 decimal places. Since the raw average was 20.0769, the final reported figure becomes **20.077**. It is important to remember that flexibility is key: to round to a different number of decimal places (e.g., two or four), simply adjust the final argument within the **ROUND** function from 3 to the desired number.

## Method 2: Rounding the Average to the Nearest Whole Integer

In contrast to highly precise reporting, many business summaries, dashboards, or presentations require averages to be presented solely as whole numbers--or the nearest integer. This simplifies communication, removing ambiguity and focusing the viewer on the overall magnitude rather than fractional components. To achieve this, the second argument of the ROUND function must be set explicitly to zero (0).

Setting the rounding precision argument to 0 signals to Google Sheets that the resulting number should have zero digits after the decimal separator, effectively rounding the calculated average to its closest whole number value following standard mathematical rounding rules (0.5 and above rounds up, below 0.5 rounds down).

The specialized formula structure for rounding the average to the nearest whole integer is:

**=ROUND(AVERAGE(A2:A14), 0)**


This command first computes the average of the range **A2:A14**, and then, using the argument **0**, ensures that the output is rounded perfectly to the nearest whole integer, providing a clean, non-fractional result for clear reporting.

### Detailed Example 2: Achieving the Nearest Integer

Using the exact same dataset illustrated previously (which yielded a raw average of 20.0769), we apply Method 2 to obtain the simplest possible representation of the mean. This example is vital for understanding how to quickly transform complex calculated values into easily understandable metrics, such as counts of people, vehicles, or discrete units.

By entering the formula `=ROUND(AVERAGE(A2:A14), 0)` into a designated output cell, we instruct the spreadsheet to perform the required aggregation and rounding. Since 20.0769 is far closer to 20 than 21, the rounding rule dictates a downward adjustment.

The following visualization confirms the implementation of this function:



D2    fx =ROUND(AVERAGE(A2:A14), 0)

	A	B	C	D
1	<b>Values</b>		<b>Average</b>	<b>Average Rounded to Integer</b>
2	14		20.07692308	20
3	19			
4	30			
5	24			
6	25			
7	25			
8	23			
9	18			
10	15			
11	13			
12	11			
13	10			
14	34			
15				
16				
17				

The execution of the formula successfully returns the average value of the cells in the range

**A2:A14**, rounded to the nearest integer. The resultant value displayed in the cell is **20**. This simple output is ideal when fractional averages are illogical or misleading in the context of the data being analyzed.

## Advanced Considerations: ROUNDUP and ROUNDDOWN

While the standard ROUND function follows typical mathematical rounding rules, there are scenarios where a user may need to enforce specific rounding behavior--always rounding up or always rounding down, regardless of the decimal value. [Google Sheets](#) accommodates these requirements through the **ROUNDUP** and **ROUNDDOWN** functions.

These functions operate with the same syntax as the standard ROUND function, taking the number (or the AVERAGE calculation) and the number of places (0 for the nearest integer) as arguments. They provide complete control over the direction of the rounding operation, which is critical in logistical planning, inventory management, or resource allocation where partial units must always be accounted for by rounding conservatively.

If you require the result to be rounded **up** to the next whole integer, even if the decimal part is small (e.g., 20.0769 becomes 21), you would substitute the ROUND function with **ROUNDUP**. Conversely, to always round **down** (e.g., 20.999 becomes 20), you use **ROUNDDOWN**. This flexibility ensures that the spreadsheet calculations align perfectly with real-world constraints and decision-making logic.

## Conclusion: Streamlining Data Reporting

The ability to calculate and simultaneously round the average of a data set using nested functions in [Google Sheets](#) is a fundamental skill for generating clear and professional reports. Whether the requirement is precise decimal control (Method 1) or simplified whole number representation (Method 2), the combination of the **AVERAGE** and **ROUND** functions offers an efficient, single-cell solution.

By leveraging these techniques, users can significantly streamline their data processing workflow, reducing manual formatting errors and accelerating the transition from raw data calculation to actionable reporting metrics. Mastery of these nested formulas allows for highly customizable data outputs tailored exactly to the needs of the audience.

The following tutorials explain how to perform other common tasks in Google Sheets:

How to calculate weighted averages using the **SUMPRODUCT** function.

Using conditional formatting based on calculated averages.

Advanced statistical analysis functions for deeper data insights.