

How can the geometric mean be calculated in R, and what are some examples of its use?

Authored by
stats writer

May 2, 2024

RECOMMENDED CITATION

stats writer (2024). *How can the geometric mean be calculated in R, and what are some examples of its use?*. PSYCHOLOGICAL SCALES. Retrieved from <https://scales.arabpsychology.com/?p=141907>

The geometric mean is a mathematical concept used to calculate the average of a set of numbers, taking into account their multiplicative relationships. In R, the geometric mean can be calculated using the "geometric.mean()" function from the Hmisc" package. This function takes a vector of numerical values as an input and returns the geometric mean of those values.

One common application of the geometric mean in R is in financial analysis, where it is used to calculate the average growth rate of a series of investments over time. Another example is in biology, where it is used to determine the average growth rate of a population over a period of time.

To calculate the geometric mean in R, the following steps can be followed:

1. Install and load the Hmisc" package.
2. Create a vector containing the numerical values for which the geometric mean needs to be calculated.
3. Use the "geometric.mean()" function, passing the vector as an argument.
4. The function will return the geometric mean of the values.

In summary, the geometric mean is a useful tool in R for calculating the average of a set of numbers, especially when the numbers have multiplicative relationships. Its applications can be seen in various fields such as finance, biology, and other areas where the growth rate needs to be determined.

Calculate Geometric Mean in R (With Examples)

You can use the following syntax to calculate the geometric mean of a set of numbers in R:

```
exp(mean(log(x)))
```

The following examples show how to use this function in practice.

Example 1: Calculate Geometric Mean of Vector

The following code shows how to calculate the geometric mean for a single vector in R:

```
#define vector
```

```
x <- c(4, 8, 9, 9, 12, 14, 17)
```

```
#calculate geometric mean of values in vector
```

```
exp(mean(log(x)))
```

```
9.579479
```

Example 2: Calculate Geometric Mean of Vector with Zeros

If your vector contains zeros or negative numbers, the formula above will return a 0 or a NaN.

To ignore zeros and negative numbers when calculating the geometric mean, you can use the following formula:

```
#define vector with some zeros and negative numbers
```

```
x <- c(4, 8, 9, 9, 12, 14, 17, 0, -4)
```

```
#calculate geometric mean of values in vector
```

```
exp(mean(log(x)))
```

```
9.579479
```

Example 3: Calculate Geometric Mean of Columns in Data Frame

The following code shows how to calculate the geometric mean of a column in a data frame:

```
#define data frame
```

```
df <- data.frame(a=c(1, 3, 4, 6, 8, 8, 9),
```

```
b=c(7, 8, 8, 7, 13, 14, 16),
```

```
c=c(11, 13, 13, 18, 19, 19, 22),
```

```
d=c(4, 8, 9, 9, 12, 14, 17))
```

```
#calculate geometric mean of values in column 'a'
```

```
exp(mean(log(df$a)))
```

```
4.567508
```

And the following code shows how to calculate the geometric mean of multiple columns in a data frame:

```
#define data frame
```

```
df <- data.frame(a=c(1, 3, 4, 6, 8, 8, 9),
```

```
b=c(7, 8, 8, 7, 13, 14, 16),
```

```
c=c(11, 13, 13, 18, 19, 19, 22),
```

```
d=c(4, 8, 9, 9, 12, 14, 17))
```

```
#calculate geometric mean of values in column 'a', 'b',
```

and 'd'

apply(df, 2, function(x) exp(mean(log(x))))

a b d

4.567508 9.871128 9.579479

ARABPSYCHOLOGY.COM