

How to Run a Correlation Test in SPSS: A Step-by-Step Guide

Authored by
stats writer

January 23, 2026

RECOMMENDED CITATION

stats writer (2026). *How to Run a Correlation Test in SPSS: A Step-by-Step Guide*.

PSYCHOLOGICAL SCALES. Retrieved from <https://scales.arabpsychology.com/?p=127249>

A correlation test is a fundamental statistical method used across numerous disciplines, providing insight into the nature of the linear relationship between two quantitative variables. This analysis determines both the strength (how closely the points align to a line) and the direction (positive or negative) of that relationship. For researchers utilizing statistical software, performing this analysis efficiently is crucial. This detailed guide outlines the precise procedures necessary to conduct a powerful Pearson correlation coefficient test using SPSS Statistics.

In order to perform a correlation test in SPSS, the following robust steps can be followed, ensuring accurate data preparation, execution, and meaningful interpretation of the resulting statistics.

Perform a Correlation Test in SPSS

The Role of the Pearson Correlation Coefficient

In quantitative statistics, the most frequently employed measure for assessing the linear association between two continuous variables is the Pearson correlation coefficient (r). Before running the test in SPSS, it is essential to understand that the goal of the correlation test is not just to calculate this coefficient, but to determine whether the resulting association is statistically significant--meaning it is unlikely to have occurred by random chance.

To determine statistical significance, we calculate a t-score and a corresponding p-value. This process relies upon a formal framework of null and alternative hypotheses. The hypotheses structure guides the interpretation of the output, specifically focusing on whether we can reject the assumption of no correlation.

Understanding the Null and Alternative Hypotheses

A correlation test operates under two competing hypotheses which define the scope of the statistical inference. These hypotheses frame the question of whether the observed relationship holds true for the larger population from which the sample data was drawn. It is critical to grasp these concepts before executing the analysis in SPSS.

The standard hypotheses utilized for assessing correlation are:

H₀: The correlation between the two variables in the population **is not** statistically significant (i.e., the population correlation coefficient, denoted as ρ , equals zero). This is the assumption of no effect or no linear relationship.

H_A: The correlation between the two variables in the population **is** statistically significant (i.e., the population correlation coefficient, ρ , is not equal to zero). This hypothesis asserts that a meaningful linear relationship exists.

The decision to reject or fail to reject the null hypothesis (H_0) rests entirely on the resulting significance level (p-value) generated by SPSS. If this p-value falls below a predefined threshold (typically the alpha level, such as $\alpha = .05$), we possess sufficient statistical evidence to reject H_0 and conclude that the relationship is statistically significant.

Step-by-Step Guide: Performing the Bivariate Correlation Test in SPSS

The process of running a correlation test in SPSS is streamlined through the menu interface, specifically utilizing the Bivariate correlation function. This method allows users to quickly calculate the relationship between pairs of variables. Following these steps ensures accurate execution of the analysis:

Data Preparation and Import: Begin by launching the SPSS software and import the data set that contains the two variables of interest. Ensure your data is clean and meets the assumptions for Pearson correlation (e.g., continuous data and approximate normality).

Accessing the Analysis Menu: Navigate to the main menu bar located at the top of the SPSS window. Select the **Analyze** option. From the dropdown menu that appears, choose **Correlate**, followed by **Bivariate**.

Variable Selection: A new dialog box titled "Bivariate Correlations" will open. Select the two specific variables you wish to test for correlation from the list on the left. Move these variables to the **Variables** box on the right by clicking on the arrow button.

Setting Correlation Type and Parameters: Ensure that the box is checked next to **Pearson** under the list of **Correlation Coefficients**. This confirms you are using the appropriate metric for linear correlation between continuous data. The default significance test is typically sufficient.

Optional Statistics: If you require supplementary descriptive statistics alongside your correlation output, click the **Statistics** tab. Here, you can choose to include additional statistics, such as means and standard deviations, which are helpful for data exploration.

Executing the Test: Once all parameters are verified and the correct variables are selected, click **OK** to run the correlation test. The results will be displayed immediately in the SPSS Output Viewer window.


Illustrative Example: Calculating Correlation in SPSS

To demonstrate this procedure clearly, consider a practical scenario where we have two variables, labeled **X** and **Y**, entered into our SPSS data file. Our objective is to determine the strength and significance of the linear association between these two variables.


Suppose the initial data setup in SPSS appears as follows, illustrating the values recorded for variables X and Y:

	 X	 Y	var	var
1	4	22		
2	6	25		
3	7	24		
4	7	20		
5	8	26		
6	10	24		
7	11	35		
8	12	30		
9	14	33		
10	15	37		
11	15	36		
12	17	26		
13	19	35		
14	20	34		
15	21	28		
16				
17				
18				
19				

We proceed by navigating through the menu commands detailed previously: Click the **Analyze** tab, then select **Correlate**, and finally choose **Bivariate**. This action opens the necessary configuration dialog box.



Within the "Bivariate Correlations" window, we drag both the **X** and **Y** variables into the **Variables** selection box. We confirm that **Pearson** is selected as the desired correlation coefficient type.



After verifying all settings, we click **OK**. The output shows a correlation matrix between X and Y, summarizing the results of our correlation test.

➔ **Correlations**

Correlations

		X	Y
X	Pearson Correlation	1	.651**
	Sig. (2-tailed)		.009
	N	15	15
Y	Pearson Correlation	.651**	1
	Sig. (2-tailed)	.009	
	N	15	15

** . Correlation is significant at the 0.01 level (2-tailed).

Interpreting the Correlation Coefficient (r) and Relationship Direction

The primary result found in the output viewer is the correlation coefficient (r), which ranges from -1 to 1. This value immediately informs us about the strength and direction of the linear association between the two variables. Stronger relationships are indicated by coefficients closer to 1 or -1.

From the example output, we can observe the following key values:

Pearson correlation coefficient (r): **.651**

p-value of Pearson correlation coefficient: **.009**

The positive value of **.651** indicates a **positive correlation**. This means that as one variable increases, the other also tends to increase. Conversely, a negative value would indicate a negative correlation, where an increase in one variable is associated with a decrease in the other.

Note: Since the Pearson correlation coefficient (**.651**) was a positive value, this indicates that there is a positive relationship between the two variables.

Analyzing Statistical Significance (The P-Value)

The significance level (p-value) is crucial for determining if the correlation observed in the sample data is reliable enough to generalize to the larger population. It indicates the probability of obtaining the calculated correlation coefficient by pure chance, assuming no actual relationship exists (H_0 is true).

As previously established, we rely on the following decision rule:

If the p-value is less than 0.05 (or the chosen alpha level), we reject H_0 .

If the p-value is greater than 0.05, we fail to reject H_0 .

Since the p-value in the output (**.009**) is significantly less than the standard threshold of .05, we reject the null hypothesis. We have sufficient statistical evidence to conclude that the correlation between variables X and Y is statistically significant, validating the observed positive linear relationship.

Further Exploration of Correlation Coefficients

Mastering the correlation test in SPSS is an essential skill for any quantitative analyst. While the Pearson method is the standard for continuous data, SPSS also offers options for Spearman's rho or Kendall's tau for non-parametric or ordinal data. Selecting the correct coefficient based on your data type is key to obtaining valid results.

The following tutorials provide additional information about correlation coefficients:

[An Introduction to the Pearson Correlation Coefficient](#)

ARABPSYCHOLOGY.COM