How to Perform a Two-Sample t-Test on Your TI-84 Calculator to Find Significant Differences

Authored by

stats writer

March 11, 2026

ARABPSYCHOLOGY.COM

RECOMMENDED CITATION

stats writer (2026). *How to Perform a Two-Sample t-Test on Your TI-84 Calculator to Find Significant Differences*. PSYCHOLOGICAL SCALES. Retrieved from https://scales.arabpsychology.com/?p=135170

Foundations of the Two Sample t-test

The **Two Sample t-test** is a fundamental statistical procedure used to determine if there is a significant difference between the **means** of two **independent** groups. Unlike a one-sample test that compares a single group to a known value, this test evaluates two distinct populations, such as a control group versus a treatment group. The core objective is to discern whether the observed variations in **average** scores are statistically meaningful or simply the result of random sampling error. This makes the test invaluable in fields ranging from medicine to engineering, where comparing outcomes is essential for decision-making.

In practice, the **Two Sample t-test** relies on the construction of a **null hypothesis**, which typically posits that there is no difference between the population means. Researchers then gather sample data and calculate a **t-statistic** to see how far the sample results deviate from the null hypothesis. If the resulting **p-value** is sufficiently low, the null hypothesis is rejected in favor of the **alternative hypothesis**. This transition from raw data to statistical inference requires precision, which is why utilizing a reliable tool like the **TI-84** calculator is so common in both academic and professional settings.

Understanding the mathematical framework of the test is crucial before performing the calculation. The test assumes that the data points within each group are collected independently and that the underlying populations follow a **normal distribution**. While the **t-test** is remarkably robust against minor deviations from normality, especially with larger sample sizes, it is always wise to perform preliminary data visualization. By leveraging the computational power of modern graphing calculators, users can bypass manual arithmetic and focus on the interpretation of the **statistical significance** of their findings.

Assessing Assumptions for Validity

Before proceeding with a **Two Sample t-test** on your **TI-84**, it is imperative to verify that your data meets the necessary statistical assumptions. The first major assumption is **independence**; the observations in the first sample must not have any influence on the observations in the second sample. For instance, if you are testing the height of students in two different schools, the selection of a student in School A should not affect the selection of a student in School B. If the samples are related, such as measuring the same person before and after a treatment, a paired t-test would be required instead.

The second assumption involves the distribution of the data, specifically the **normal distribution**. For the results of a **t-test** to be valid, the populations from which the samples are drawn should ideally be bell-shaped. However, thanks to the **Central Limit Theorem**, the test remains effective even if the data is not perfectly normal, provided the sample size is sufficiently large (typically n >

ARABPSYCHOLOGY.COM

30). When working with smaller samples, as seen in many experimental trials, it is recommended to check for outliers or extreme skewness that could distort the **mean** and **standard deviation**.

Lastly, one must consider the **variance** within each group. Classical versions of the t-test assume that both populations have equal variances, a condition known as **homoscedasticity**. If the variances are significantly different, the standard t-test may produce misleading results. Fortunately, the **TI-84** allows users to choose between **pooled** and non-pooled methods, with the latter utilizing **Welch's t-test** to account for unequal variances. This flexibility ensures that the **t-statistic** and **degrees of freedom** are calculated with the highest possible accuracy for the specific data set at hand.

Capabilities of the TI-84 Plus Series

The **TI-84 Plus** series is widely recognized as the industry standard for educational and introductory statistical analysis. Its robust internal software includes a comprehensive suite of "TESTS" designed to handle everything from simple proportions to complex **ANOVA**. For students and researchers, the ability to perform a **Two Sample t-test** directly on the device eliminates the need for complex software or tedious manual calculations involving **t-distribution** tables. The calculator's interface is designed to prompt the user for specific inputs, reducing the likelihood of procedural errors during the testing phase.

Beyond simple calculation, the **TI-84** provides a visual context for statistics. Users can plot their data using histograms or box-and-whisker plots to check for **normality** and outliers before running the test. This integration of graphical and numerical analysis fosters a deeper understanding of how **standard deviation** and sample size influence the final **p-value**. Whether you are using the classic **TI-84 Plus** or the modern **TI-84 Plus CE** with its high-resolution color screen, the logic and menu structures remain consistent, making it a versatile tool for long-term use.

One of the most powerful features of the **TI-84** is its dual-input mode for statistical tests. Users can choose to input "Data," which involves entering raw values into lists (L1, L2), or "Stats," which allows for the entry of pre-calculated summary statistics like the **mean** and **Sx**. This flexibility is particularly useful when you are performing an analysis based on data found in a textbook or a research paper where the raw observations are not provided. By mastering these menus, you can quickly pivot between different types of **hypothesis testing** with confidence and speed.

Preparing for Calculation: Data Entry and Lists

To begin your analysis, you must first organize your data. If you have raw data points for both groups, the most efficient method is to use the **List** editor on your **TI-84**. Press the **STAT** button and select **1:Edit...** to open the list interface. Typically, researchers will enter the data for the first sample into **L1** and the data for the second sample into **L2**. Ensure that each data point is entered

ARABPSYCHOLOGY.COM

accurately, as a single typo can significantly alter the **sample mean** and the resulting **t-statistic**.

If you are not using raw data but instead have summary statistics--such as the **mean**, **standard deviation**, and sample size--you can skip the list entry phase entirely. This is common in exam scenarios where the problem provides the summary values directly. Regardless of the input method, it is helpful to clearly label your groups on paper to avoid confusing the "Control" parameters with the "Treatment" parameters. Proper organization at this stage streamlines the **Two Sample t-test** process and ensures that the alternative hypothesis is set up correctly in the following steps.

Once the data is ready, double-check the sample sizes (**n**). The **Two Sample t-test** does not require the groups to have an equal number of observations, but the sizes should be recorded accurately. Small sample sizes increase the **standard error** and make it more difficult to find **statistical significance** unless the difference between the means is very large. With your data organized and your calculator powered on, you are now ready to navigate the specialized "TESTS" menu to perform the actual **2-SampTTest**.

Accessing the 2-SampTTest Menu

The core of the statistical functionality on the **TI-84** is housed within the **STAT** menu. To perform a **Two Sample t-test**, begin by pressing the **STAT** key. Use the right arrow key to scroll over to the **TESTS** tab. This menu contains a variety of procedures for **hypothesis testing** and **confidence intervals**. Scroll down using the down arrow key until you find **4:2-SampTTest...**. It is important not to confuse this with "2-SampZTest," which is only used when the population standard deviations are known--a rare occurrence in real-world research.

After highlighting **2-SampTTest**, press **ENTER**. This will open a new screen where you will define the parameters of your test. The first choice you must make is the **Inpt** (Input) type. If you have entered raw data into lists, highlight **Data** and press **ENTER**. If you have the **mean** and **standard deviation** ready, highlight **Stats** and press **ENTER**. Choosing the correct input method is vital, as it changes the fields that the calculator displays for the subsequent steps.

ARABPSYCHOLOGY.COM



As shown in the interface, the **TI-84** provides a clean layout for entering your variables. If you chose **Data**, the calculator will ask for the list names (List1 and List2) and the frequency (Freq1 and Freq2), which should usually be set to 1. If you chose **Stats**, you will see fields for the **average** of each group, their **standard deviations**, and their sample sizes. Taking a moment to ensure you are in the correct menu prevents the frustration of having to re-enter data later in the process.

Inputting Summary Statistics (Stats Mode)

When working in **Stats** mode, the **TI-84** requires six specific values to perform the **Two Sample t-test**. First, you will enter **x1**, which represents the **mean** of the first sample. Following this, you must enter **Sx1**, the **sample standard deviation** for that same group. It is important to use the sample standard deviation (Sx) rather than the population standard deviation (σx) provided by some calculations, as the **t-test** is specifically designed for cases where population parameters are unknown. Finally, enter **n1**, the total number of observations in the first group.

Repeat this process for the second group by entering values for **x2**, **Sx2**, and **n2**. Accuracy here is paramount. For example, if you are comparing fuel treatments, **x1** might be the **mean** mpg of the control cars, while **x2** is the **mean** mpg of the cars receiving the new treatment. Because the **TI-84** uses these values to calculate the **t-statistic**, even a small decimal error can lead to a different **p-value** and potentially change the conclusion of your entire study. Always double-verify your entries against your source data before moving to the hypothesis selection.

In addition to these numerical inputs, the calculator allows you to define the nature of your **statistical hypothesis**. This is where you specify whether you are looking for any difference at all, or if you are specifically testing if one group is greater than or less than the other. The **TI-84** handles the complex integration of the **t-distribution** curve behind the scenes, but it relies on you to provide these correct starting values. Once these fields are filled, you are ready to address the logical direction of your test.

ARABPSYCHOLOGY.COM

Defining the Alternative Hypothesis

Choosing the correct **alternative hypothesis** is one of the most critical steps in **hypothesis testing**. On the **TI-84**, you will see a row labeled **μ1** with three options: **? μ2**, **<μ2**, and **>μ2**. The **? μ2** option is used for a **two-tailed test**, which checks for any significant difference between the two groups, regardless of which one is larger. This is the most conservative and common approach in scientific research, as it does not assume the direction of the effect before the data is analyzed.

If you have a specific prediction--for example, that a new medicine will definitely increase recovery rates--you might choose a **one-tailed test** (either **<μ2** or **>μ2**). However, one-tailed tests are generally scrutinized more heavily because they make it easier to achieve **statistical significance** in one direction while completely ignoring potential differences in the other. In most academic settings, unless specified otherwise, the **two-tailed** (?) option is the standard choice. Highlighting your selection and pressing **ENTER** tells the **TI-84** which area of the **t-distribution** to calculate for the **p-value**.

The selection you make here directly impacts the **p-value**. For a **two-tailed test**, the calculator doubles the area in the tail of the **t-statistic** to account for differences in both directions. This level of detail is what makes the **TI-84** such a powerful tool for students; it automates the most complex part of the **Two Sample t-test** while requiring the user to understand the underlying logic of their research question. Once the hypothesis is set, you will reach the final toggle: the choice to pool your data or not.

The Nuance of Pooled Variance

The **Pooled** option on the **TI-84** asks whether you want to combine the **variances** of the two samples into a single estimate. This is only statistically appropriate if you have a reason to believe that the two populations have equal **standard deviations**. In many traditional statistics courses, pooling was common because it simplified manual calculations. However, in modern practice, **Pooled: No** is the recommended default. Selecting "No" performs **Welch's t-test**, which is more robust and accurate when the variances or sample sizes are unequal.

When you select **Pooled: No**, the calculator uses a more complex formula to determine the **degrees of freedom**. This often results in a non-integer value for **df**, which you will see in the final results. This adjustment is necessary to ensure that the **p-value** is not artificially deflated, which would increase the risk of a **Type I error** (a false positive). By choosing not to pool, you are allowing the **TI-84** to provide a more reliable and honest assessment of the **statistical significance** of your data.

After making your selection for **Pooled**, move the cursor down to **Calculate** and press **ENTER**. Alternatively, you can select **Draw**, which will show you the **t-distribution** curve with the shaded

ARABPSYCHOLOGY.COM

area representing the **p-value**. Most users prefer **Calculate** because it provides a clear list of all the numerical values needed for a formal report. The calculator then processes the inputs and displays the results screen, which contains the final verdict of your **Two Sample t-test**.



Analyzing the Calculator Output

Once you hit calculate, the **TI-84** presents a screen filled with data. The most important value for most researchers is the **p-value** (labeled as **p**). This number represents the probability of observing a difference as large as the one in your samples if the **null hypothesis** were actually true. If **p** is less than your **significance level** (usually 0.05), you have sufficient evidence to reject the null hypothesis and conclude that a significant difference exists between the **means**.

The output also displays the **t-statistic** (labeled as **t**), which indicates how many standard errors the difference between the **sample means** is from zero. A larger absolute value for **t** generally corresponds to a smaller **p-value**. Additionally, the **degrees of freedom** (**df**) are shown. As mentioned previously, if you did not pool the variances, this number will likely be a decimal. This is the **Satterthwaite approximation** used in **Welch's t-test** to provide a more accurate result when group variances differ.



The rest of the output screen lists the values you entered--such as **x1**, **x2**, **Sx1**, **Sx2**, **n1**, and **n2**. This allows you to double-check that you didn't make an entry error. If the results look unexpected,

ARABPSYCHOLOGY.COM

checking these values is the first step in troubleshooting. Interpreting these results correctly is the final piece of the **Two Sample t-test** puzzle, allowing you to turn raw numbers into a meaningful conclusion about your research topic.

Practical Application: A Fuel Efficiency Case Study

To illustrate the process, consider a study where researchers test a new fuel treatment to see if it improves **average** mpg. They use two groups of 12 cars each. The control group (no treatment) shows a **mean** of 21 mpg with a **standard deviation** of 2.73. The treatment group shows a **mean** of 22.75 mpg with a **standard deviation** of 3.25. By entering these values into the **2-SampTTest** menu on the **TI-84** and selecting a **two-tailed test** (μ1? μ2) without pooling, we can determine the **statistical significance** of the fuel treatment.

Upon calculation, the **TI-84** provides a **t-statistic** of approximately -1.428 and a **p-value** of 0.1677. Since this **p-value** is significantly higher than the standard 0.05 threshold, we fail to reject the **null hypothesis**. In plain language, this means that although the treatment group had a slightly higher **average** mpg, the difference was not large enough to rule out random chance. The experiment does not provide enough evidence to say the fuel treatment actually works for this specific car model.

This example highlights the importance of the **Two Sample t-test** in preventing false claims. Without the rigors of **hypothesis testing**, one might look at the 22.75 versus 21.00 mpg and conclude the treatment is effective. However, the **TI-84** analysis reveals that the **variance** within the groups is too high and the sample size is too small to be certain. By following these structured steps on your calculator, you can ensure that your own data analysis is held to the same high standard of scientific integrity.

ARABPSYCHOLOGY.COM