

How to Calculate the Average of Cells Containing Numbers in Excel

Authored by
stats writer

November 28, 2025

RECOMMENDED CITATION

stats writer (2025). *How to Calculate the Average of Cells Containing Numbers in Excel*. PSYCHOLOGICAL SCALES. Retrieved from <https://scales.arabpsychology.com/?p=100832>

When working with complex datasets in Excel, calculating the arithmetic mean of a range is typically achieved using the built-in AVERAGE function. This powerful function efficiently processes a specified range of cells and ignores any non-numeric entries, returning the mean of the numerical values present. However, data analysis often requires a more sophisticated condition: calculating the average of values in one column only if a corresponding cell in a separate criteria column meets a specific standard, such as strictly containing a number. This necessity arises frequently when dealing with mixed data types where identifiers might be numerical or alphanumeric, but calculations must only proceed if the identifier is confirmed as purely numerical.

Standard conditional functions, such as **AVERAGEIF** or **AVERAGEIFS**, are excellent for basic numerical or textual comparisons within a single range or between two ranges based on equality. Nevertheless, they lack the inherent capability to perform a data type check (e.g., "Is this cell a number?") and then use that check to filter values in an adjacent column. To achieve this level of precise conditional calculation, we must employ an advanced array formula structure, leveraging the efficiency of the **SUMPRODUCT** function combined with the logic provided by the **ISNUMBER** function. This approach allows us to create a dynamic filter mask that ensures only qualifying records are included in both the sum and the count components of the average calculation.

The Logic Behind Conditional Data Type Checking

The foundation of this conditional average calculation rests on establishing a Boolean mask--a series of TRUE or FALSE values--that identifies precisely which cells in the criteria range are numerical. This mask acts as a gatekeeper, determining which values from the data range will be included in the final calculation. Since Excel formulas require numerical inputs for arithmetic operations, the TRUE/FALSE results must be coerced into their numerical counterparts: 1 for TRUE and 0 for FALSE.

The key functions enabling this specialized filtering are the ISNUMBER function, which performs the type check, and the double unary operator (--), which handles the coercion. When this resulting array of 1s and 0s is multiplied against the corresponding values we wish to average, any value paired with a 0 is nullified, and any value paired with a 1 is retained. This crucial multiplication step effectively isolates the data points that meet the condition.

Furthermore, we must ensure that the denominator in our averaging calculation reflects the count of only the qualifying records, not the total number of records in the value range. If the resulting sum of qualified values is divided by the total count of the value range, the result will be incorrectly low. This requirement is satisfied elegantly by using the standard COUNT function applied specifically to the criteria range, as **COUNT** inherently tallies only numerical entries, perfectly matching our criteria.

Introducing the Robust Conditional Averaging Formula

The structure below represents the most efficient and robust method for calculating the average of values in Range B, contingent upon the corresponding cells in Range A being numerical. This formula utilizes array processing capabilities inherent in the SUMPRODUCT function to handle the conditional summation, paired with the COUNT function for the denominator.

You can use the following formula to calculate the average in Excel only for the cells that contain a number in a corresponding range:

=SUMPRODUCT(--ISNUMBER(A2:A11), B2:B11) / COUNT(A2:A11)

This particular formula will calculate the average of the values located in the range **B2:B11**, but only for those rows where the corresponding cell in the criteria range **A2:A11** contains a recognized numerical value. If either of the ranges is empty, contains errors, or if the ranges are not of the exact same size, the formula may return unexpected results or an error, emphasizing the need for carefully structured data input. Understanding the role of each component within this sophisticated structure is essential for accurate application across various datasets.

Deconstructing the Formula's Numerator: Conditional Summation

The numerator, `SUMPRODUCT(--ISNUMBER(A2:A11), B2:B11)`, is responsible for performing the conditional aggregation. The SUMPRODUCT function is chosen because it inherently handles array operations efficiently, multiplying corresponding elements of arrays and summing the resultant products without requiring the user to enter the formula using Ctrl+Shift+Enter (a requirement for older array formulas).

The operation begins with the ISNUMBER function evaluating the criteria range **A2:A11**. This step generates a memory array consisting entirely of TRUE or FALSE values, depending on the content of each cell. TRUE indicates a number is present; FALSE indicates text, blanks, or other non-numeric data.

Next, the double unary operator (`--`) acts upon this array. This mechanism is crucial as it coerces the Boolean array into an array of integers (1s and 0s). This resulting array serves as the masking filter. Finally, SUMPRODUCT takes this mask array and multiplies it element-by-element by the values in the data range **B2:B11**. Only the values in **B2:B11** corresponding to a '1' in the mask are retained for the final sum. Values corresponding to a '0' are multiplied by zero, effectively excluding them from the total. This process ensures the numerator contains the sum of only the conditionally qualifying records.

The Denominator: Accurate Conditional Counting

A correct average requires dividing the conditional sum by the accurate count of records that met the condition. The denominator, `COUNT(A2:A11)`, solves this requirement with remarkable simplicity.

The standard `COUNT` function, when applied to a range, is designed specifically to count only the cells within that range that contain numerical data. Since our exact condition is "does the cell contain a number," applying `COUNT` directly to the criteria range **A2:A11** yields the precise count of rows that qualify for inclusion in the average. This avoids the need for a redundant `SUMPRODUCT` calculation for the denominator, streamlining the formula and improving performance. If the criteria were different--for example, counting cells equal to "Active"--we would instead use a conditional count like `SUMPRODUCT(--(A2:A11="Active"))`. However, for number checking, `COUNT(A2:A11)` is the optimized solution.

Practical Example Setup: Analyzing Sales Performance

To solidify the understanding of this technique, let us apply it to a practical data analysis scenario. Imagine a business tracking employee sales performance. Employee identification (ID) is recorded in Column A, and their total sales are recorded in Column B. Due to varied employee statuses (permanent staff, temporary contractors, etc.), the ID column contains a mixture of true numerical IDs and text-based identifiers (such as "N/A" or unique alphanumeric codes). Our objective is to calculate the average sales performance, focusing strictly on employees identified by a purely numeric ID, thus excluding all employees with text-based identifiers.

The dataset below illustrates this typical mixed-data structure:

	A	B	C	D	E
1	ID	Sales			
2	A	22			
3	B	14			
4	C	29			
5	4	19			
6	D	31			
7	E	10			
8	7.11	4			
9	8	8			
10	H	12			
11	10	17			
12					
13					
14					
15					
16					
17					
18					
19					
20					
21					

In this sample data, the range A2:A11 holds the ID status, which includes numerical values, text strings, and blank cells. The corresponding sales figures in B2:B11 must be conditionally averaged. We need to identify the sales figures associated with rows where A2:A11 is numeric and average only those specific values.

Implementing and Visualizing the Conditional Formula

We will now implement the derived formula to solve the conditional averaging problem based on the sales data provided. This demonstration reinforces how the ISNUMBER function and SUMPRODUCT function collaborate to perform the necessary screening and aggregation.


The formula used to calculate the average sales made exclusively by employees whose ID is a numerical value in range A2:A11 is:

=SUMPRODUCT(--ISNUMBER(A2:A11), B2:B11) / COUNT(A2:A11)

When this formula is entered into a target cell, Excel processes the array operations, returning a single, calculated average. The result represents the arithmetic mean of the sales figures that

passed the numerical ID check.

The following screenshot shows how the formula is applied and the resulting calculated value:



The screenshot shows an Excel spreadsheet with the following data:

	A	B	C	D	E	F	G	H	I
1	ID	Sales		Avg. Sales for Numeric ID's					
2	A	22		12					
3	B	14							
4	C	29							
5	4	19							
6	D	31							
7	E	10							
8	7.11	4							
9	8	8							
10	H	12							
11	10	17							
12									
13									
14									
15									
16									
17									
18									
19									
20									
21									

The calculated average sales among employees with a numeric ID is determined to be exactly **12**. This value successfully filters out all sales records corresponding to non-numeric IDs, providing a clean average for the targeted subset of the workforce.

Manual Verification of the Calculated Average

To confirm the accuracy and methodology of the array formula, it is prudent to manually identify the qualifying data points and perform the calculation independently. This process validates that the formula correctly identified the numerical criteria and excluded all non-numerical entries.

We manually identify the rows where the ID column (A) contains a numeric value. These are the records where the ISNUMBER function returns TRUE:

	A	B	C	D	E	F
1	ID	Sales		Avg. Sales for Numeric ID's		
2	A	22		12		
3	B	14				
4	C	29				
5	4	19				
6	D	31				
7	E	10				
8	7.11	4				
9	8	8				
10	H	12				
11	10	17				
12						
13						
14						
15						
16						
17						
18						

The qualifying sales figures corresponding to the numerical IDs are 19 (Row 2), 4 (Row 4), 8 (Row 6), and 17 (Row 7). The total number of qualifying records is 4.

The average sales made by these selected employees is calculated as:

$$\text{Average Sales} = (19 + 4 + 8 + 17) / 4 = 48 / 4 = 12.$$

This manual verification confirms that the conditional calculation using the **SUMPRODUCT/ISNUMBER/COUNT** formula is entirely accurate, providing a powerful, dynamic tool for data type specific conditional averaging in complex Excel models.

Summary of Advantages and Advanced Considerations

The primary advantage of using the SUMPRODUCT function for this task lies in its ability to handle array arithmetic without the need for manual array entry, making the formula easier to deploy and modify. It provides a highly portable solution that works across most versions of Excel.

While modern versions of Excel (such as Microsoft 365) offer dynamic array alternatives, potentially using the **FILTER** function in combination with the AVERAGE function (e.g., `AVERAGE(FILTER(B2:B11, ISNUMBER(A2:A11)))`), the **SUMPRODUCT** method remains an essential technique for analysts supporting users on older software versions or when performance needs to be optimized for very large, highly calculated spreadsheets. The intelligent use of the

COUNT function applied to the criteria range provides the cleanest method for generating the denominator, specifically tailored to the "is number" condition, solidifying this formula as the optimal solution for conditional averaging based on data type.

ARABPSYCHOLOGY.COM