

How to Easily Calculate Quintiles in Excel

Authored by
stats writer

December 6, 2025

RECOMMENDED CITATION

stats writer (2025). *How to Easily Calculate Quintiles in Excel*. PSYCHOLOGICAL SCALES.
Retrieved from <https://scales.arabpsychology.com/?p=106075>

Quintiles are a fundamental type of statistical measure used for comprehensive data analysis, designed to divide a data set into five perfectly equal parts based on frequency. Understanding how to calculate these measures is essential for researchers and analysts working with distribution data. While some statistical platforms use dedicated functions, in Excel, the calculation relies primarily on the powerful PERCENTILE function, though the outdated QUARTILE function was sometimes referenced in legacy guides. The key difference is that quintiles require dividing the data into 20% increments, necessitating the use of percentile-based calculations (20%, 40%, 60%, 80%). This guide provides a detailed, step-by-step approach to calculating and interpreting these critical measures directly within your spreadsheet software.

Introduction to Quintiles and Distribution Analysis

In the realm of descriptive statistics, understanding data distribution is paramount. **Quintiles** are specific types of quantiles--numerical values that partition a ranked set of data into groups of equal size. Specifically, they split a dataset into five groups, where each group contains 20% of the total observations. This division allows analysts to quickly grasp how data points are spread across the distribution, identifying concentrations of values in the lower, middle, and upper fifths of the data. This technique is widely employed in fields like economics, finance, and social sciences to analyze income distribution, test scores, or product performance tiers.

The definition of a quintile point is based on the cumulative frequency. The first quintile (Q1) marks the boundary where 20% of all data values lie below it, and 80% lie above. Moving sequentially, the second quintile (Q2) corresponds to the 40th percentile, meaning 40% of the observations fall below this point. This progression continues until the fourth quintile (Q4), which represents the 80th percentile. The fifth quintile is, by definition, the maximum value of the dataset, representing the 100th percentile. These partition points are invaluable tools for comparative analysis, especially when evaluating data sets for skewness or inequality.

Understanding Percentiles and Quantiles

Before diving into the practical application in Excel, it is essential to solidify the relationship between quintiles and the broader concept of percentiles. A percentile divides data into 100 equal parts, making it the most granular form of quantile measure. Quintiles are simply specific percentiles: 20th, 40th, 60th, and 80th. This relationship is precisely why we must utilize the PERCENTILE function in Excel, rather than functions dedicated to quartiles (which divide data into four parts).

The use of the PERCENTILE function provides the necessary flexibility to specify the required fraction (or percentage) of the data distribution we wish to isolate. Since quintiles require fractions of 0.2, 0.4, 0.6, and 0.8, this function is perfectly suited to the task. It requires the user to input the

range of data being analyzed and the specific fraction corresponding to the desired quintile. This direct translation from percentile fraction to quintile marker streamlines the calculation process significantly.

Why Use Quintiles in Statistical Analysis?

While quartiles are perhaps the most common quantile measure used in basic statistical reporting, quintiles offer a finer, yet still easily digestible, division of the data. They are particularly useful when researchers need to categorize populations or performance metrics into five distinct, rank-ordered segments, often referred to as "fifths." This level of segmentation can reveal subtle patterns or disparities that might be obscured when using only four divisions.

Consider, for instance, analyzing student test scores or household incomes. Dividing these data sets into quintiles allows policy makers or educators to easily identify the bottom 20% (the lowest fifth), the middle 20% (the third fifth), and the top 20% (the highest fifth). This clear segmentation facilitates targeted interventions or resource allocation. The robustness of quintile analysis makes it a cornerstone of advanced descriptive statistics, providing clear cut-off points for defining performance tiers across any given dataset.

The Excel PERCENTILE Function Explained

To perform this analysis in Excel, we rely on the primary tool for quantile calculation: the PERCENTILE function. While older versions of Excel included functions like `PERCENTILE.EXC` and `PERCENTILE.INC`, the modern `PERCENTILE` function, or its inclusive variant (`PERCENTILE.INC`), is the standard approach for this calculation. This function calculates the k-th percentile of values in a range, where 'k' is the percentile coefficient between 0 and 1.

We can use the following syntax template to calculate the desired quintiles for any given data range in Excel:

=PERCENTILE(CELL RANGE, QUINTILE)

In this structure, **CELL RANGE** refers to the array of numerical data you are analyzing (e.g., A1:A20). The **QUINTILE** argument is the crucial percentile fraction. To find the first quintile (Q1), this value must be 0.2 (20%); for the second quintile (Q2), it must be 0.4 (40%), and so forth, up to 0.8 for the fourth quintile (Q4). Mastery of this simple syntax unlocks immediate access to robust quantile analysis within the spreadsheet environment.

The following practical example demonstrates precisely how to implement this function and determine the critical partition points within a sample dataset.

Example: Calculating Quintiles for a Sample Dataset in Excel

To illustrate the methodology, let us assume we have collected a sample dataset consisting of 20 distinct observations. This dataset could represent any continuous variable, such as sales figures, response times, or component weights. Organizing this data within a single column in Excel is the first step toward successful calculation.

Suppose we have the following raw numerical data listed in column A of our spreadsheet:

	A	B	C	D	E	F	G
1	Data						
2	4						
3	5						
4	5						
5	6						
6	7						
7	8						
8	12						
9	14						
10	14						
11	16						
12	17						
13	20						
14	22						
15	23						
16	24						
17	26						
18	27						
19	30						
20	35						
21	38						
22							
23							
24							
25							

Our objective is to calculate the four defining quintile boundaries (Q1, Q2, Q3, and Q4) for this specific data range (A1:A20). This requires setting up four separate formula applications, each targeting the relevant percentile coefficient (0.2, 0.4, 0.6, and 0.8) while referencing the consistent data array. This methodical approach ensures accurate partition points are identified for the entire distribution.

Step-by-Step Implementation of the Formula

To determine the specific value that marks the boundary of each fifth of the data, we apply the PERCENTILE function sequentially. For the data range A1:A20, the implementations are as follows:

For the First Quintile (Q1, 20th percentile): The formula is **=PERCENTILE(A1:A20, 0.2)**.

For the Second Quintile (Q2, 40th percentile): The formula is **=PERCENTILE(A1:A20, 0.4)**.

For the Third Quintile (Q3, 60th percentile): The formula is **=PERCENTILE(A1:A20, 0.6)**.

For the Fourth Quintile (Q4, 80th percentile): The formula is **=PERCENTILE(A1:A20, 0.8)**.

The following image vividly demonstrates the application of these formulas in Excel, resulting in the calculated quintile values adjacent to the original data:

	A	B	C	D	E	F	G	H	I
1	Data			Quintile	Value				
2	4			0.2	6.8	=PERCENTILE(\$A\$2:\$A\$21, D2)			
3	5			0.4	14				
4	5			0.6	20.8				
5	6			0.8	26.2				
6	7								
7	8								
8	12								
9	14								
10	14								
11	16								
12	17								
13	20								
14	22								
15	23								
16	24								
17	26								
18	27								
19	30								
20	35								
21	38								
22									
23									
24									
25									

Reviewing the output, we observe the specific numerical values that demarcate the boundaries between the five equal data partitions. These values are interpolation results based on the data array, ensuring that exactly 20% of the observations fall between each sequential quintile marker,

providing a clear statistical map of the data distribution.

Interpreting the Results of Quintiles

Once the numerical values for the quintiles are calculated, their true utility lies in their interpretation. These values are not data points themselves but boundary markers. Interpreting them correctly means understanding the percentage of the data that falls below each calculated figure. This interpretation is fundamental to utilizing quintile analysis effectively in statistics and reporting.

The way to interpret the calculated quintile boundaries from the example above is as follows:

The first 20% of all data values lie below the Q1 boundary value of **6.8**.

The first 40% of all data values lie below the Q2 boundary value of **14**.

The first 60% of all data values lie below the Q3 boundary value of **20.8**.

The first 80% of all data values lie below the Q4 boundary value of **26.2**.

This interpretation allows us to confirm that 20% of the data lies between 6.8 and 14, 20% lies between 14 and 20.8, and so on. If this data represented income, for example, we could state definitively that the bottom 20% of incomes are below 6.8 units, while the top 20% are above 26.2 units.

Calculating All Quintiles Simultaneously Using Array Input

For analysts who need to calculate all four quintile boundaries frequently, Excel offers a powerful shortcut using array constants within the PERCENTILE function. Instead of entering four separate formulas, a single formula can be utilized to return an array of all four calculated values (Q1, Q2, Q3, Q4) in adjacent cells. This method significantly increases efficiency, especially when dealing with multiple large data arrays.

We can use the following advanced formula structure to calculate each quintile simultaneously:

=PERCENTILE(CELL RANGE, {0.2, 0.4, 0.6, 0.8})

When entering this formula in older versions of Excel, it must be entered as an array formula (using CTRL+SHIFT+ENTER) across four selected cells. In modern versions of Excel (such as Microsoft 365), this formula uses dynamic array functionality and will automatically spill the results into adjacent cells, making the calculation instantaneous and highly robust.

The following image shows the execution and resulting output when using this array formula technique:

	A	B	C	D	E	F	G	H
1	Data							
2	4		6.8	14	20.8	26.2		
3	5							
4	5							
5	6							
6	7							
7	8							
8	12							
9	14							
10	14							
11	16							
12	17							
13	20							
14	22							
15	23							
16	24							
17	26							
18	27							
19	30							
20	35							
21	38							
22								
23								
24								

Notice that the quintiles calculated using this efficient array method perfectly match the quintiles we calculated earlier using the individual formula entries. This consistency validates both methods and demonstrates the power and flexibility of the PERCENTILE function for advanced distributional statistics within Excel.