

How to Easily Create a Bell Curve in Google Sheets

Authored by
stats writer

December 3, 2025

RECOMMENDED CITATION

stats writer (2025). *How to Easily Create a Bell Curve in Google Sheets*. PSYCHOLOGICAL SCALES. Retrieved from <https://scales.arabpsychology.com/?p=104092>


Visualizing statistical data often requires plotting a normal distribution, commonly referred to as the **Bell Curve**. This characteristic shape provides crucial insights into data dispersion and probability. While specialized software handles this easily, you can generate a clean, precise bell curve directly within Google Sheets using its powerful built-in functions.

The core methodology relies on the NORMDIST function, which calculates the probability density for a given value based on the dataset's mean and standard deviation. By generating a series of paired data points (X and Y coordinates) and then visualizing them using a scatter chart, we can accurately construct the smooth, symmetrical curve.

This comprehensive, step-by-step tutorial guides you through the process of setting up your spreadsheet, calculating the necessary distribution values, and ultimately plotting the **bell curve**. This technique ensures that your resulting visualization is dynamic and updates automatically whenever you adjust the underlying statistical parameters.

Understanding the Normal Distribution (Bell Curve)

The term "bell curve" is universally used to describe the graphical representation of the Normal Distribution. This curve is central to statistics, representing how data points cluster around the average value. Its distinctive shape, high in the middle and tapering symmetrically toward the edges, demonstrates that most data points fall close to the mean.



To accurately render this shape in Google Sheets, we need to mathematically define the curve based on specific parameters. We will begin by establishing the fundamental statistical values that govern the data set's location and spread.

Step 1: Establishing the Core Parameters (Mean and Standard Deviation)

The initial step requires defining the two most critical parameters of any normal distribution: the

Mean (μ) and the **Standard Deviation** (σ). The mean determines the central peak of the curve, while the standard deviation dictates the spread or width of the curve.

For this example, we will set up specific cells in our Google Sheet to hold these values. It is best practice to label these cells clearly for easy reference and future modification:

Enter "Mean" in cell A2 and the desired value (e.g., 50) in cell B2.

Enter "Standard Deviation" in cell A3 and the desired value (e.g., 10) in cell B3.

These reference cells (B2 and B3) will serve as dynamic inputs for our subsequent calculations using the NORMDIST function.

	A	B	C	D
1	Mean	0		
2	Standard Dev	1		
3				
4				
5				
6				
7				
8				
9				
10				
11				
12				
13				
14				
15				
16				
17				
18				

Step 2: Generating the X-Axis Data Points (Percentile Range)

To draw a smooth curve, we need a sufficient number of data points spanning the distribution's range. Statisticians typically measure the spread of a normal distribution using standard deviation units (or Z-scores). A range covering three or four standard deviations on either side of the mean is usually sufficient to capture over 99.7% of the data.

We will create a column starting in cell A5 containing our percentile values, ranging from **-4 to 4**, in small increments (e.g., 0.1). Using a smaller increment creates more data points, resulting in a significantly smoother visual representation when plotted as a scatter chart.

This range ensures that we cover the extremes of the distribution. If we use increments of 0.1, we will generate 81 data points (from -4.0, -3.9, ... 0, ... 3.9, 4.0), providing the necessary resolution for a visually accurate **bell curve**.

	A	B	C	D
1	Mean	0		
2	Standard Dev	1		
3				
4	Percentiles			
5	-4			
6	-3.9			
7	-3.8			
8	-3.7			
9	-3.6			
10	-3.5			
11	-3.4			
12	-3.3			
13	-3.2			
14	-3.1			
15	-3			
16	-2.9			
17	-2.8			
18	-2.7			
19	-2.6			
20	-2.5			
21	-2.4			

Step 3: Calculating Absolute Data Values (X-Coordinates)

Now that we have defined the relative percentile positions (Z-scores) in Column A, we need to convert them into actual data values (X-coordinates) based on our defined mean and standard deviation. These values represent the physical range of the data set along the X-axis.

In cell B5, input the formula to calculate the specific X-coordinate corresponding to the Z-score in A5, using the standard deviation value in B3 and the mean value in B2. The formula structure should be: $= (A5 * \$B\$3) + \$B\$2$. The use of absolute references (\$B\$3 and \$B\$2) is crucial, as these statistical parameters must remain constant as the formula is copied down the column.

Apply this formula to every row corresponding to the percentile range defined in Step 2 (e.g., from cell B5 down to B85). This column now represents the input values for which we will calculate the probability density.

B5	fx	=B\$1+A5*B\$2		
	A	B	C	D
1	Mean	0		
2	Standard Dev	1		
3				
4	Percentiles	Data		
5	-4	-4		
6	-3.9	-3.9		
7	-3.8	-3.8		
8	-3.7	-3.7		
9	-3.6	-3.6		
10	-3.5	-3.5		
11	-3.4	-3.4		
12	-3.3	-3.3		
13	-3.2	-3.2		
14	-3.1	-3.1		
15	-3	-3		
16	-2.9	-2.9		
17	-2.8	-2.8		
18	-2.7	-2.7		
19	-2.6	-2.6		
20	-2.5	-2.5		
21	-2.4	-2.4		

Step 4: Calculating Probability Density Function (Y-Coordinates)

The height of the **bell curve** at any given X-coordinate is determined by the Probability Density Function (PDF). In Google Sheets, this is calculated using the specialized NORMDIST function. This function requires four arguments: the X value, the mean, the standard deviation, and a logical value indicating whether to calculate the cumulative distribution (TRUE) or the probability density (FALSE).

In cell C5, enter the following formula: `=NORMDIST(B5, B$2, B$3, FALSE)`. Here, B5 is the calculated X-value, B\$2 is the fixed mean, and B\$3 is the fixed standard deviation. We use FALSE because we are interested in the height of the curve (the density) at that specific point, not the cumulative area under the curve.

Copy this formula down the column from C5 to C85. Column C now contains the Y-coordinates corresponding to the X-coordinates in Column B, effectively mapping out the precise shape of the normal distribution.

C5	Σ	=NORM.DIST(B5, \$B\$1, \$B\$2, FALSE)		
	A	B	C	D
1	Mean	0		
2	Standard Dev	1		
3				
4	Percentiles	Data	PDF	
5	-4	-4	0.000133830225	
6	-3.9	-3.9	0.000198655471	
7	-3.8	-3.8	0.000291946925	
8	-3.7	-3.7	0.000424780270	
9	-3.6	-3.6	0.000611901930	
10	-3.5	-3.5	0.000872682695	
11	-3.4	-3.4	0.001232219168	
12	-3.3	-3.3	0.001722568939	
13	-3.2	-3.2	0.002384088201	
14	-3.1	-3.1	0.003266819056	
15	-3	-3	0.004431848412	
16	-2.9	-2.9	0.00595253242	
17	-2.8	-2.8	0.007915451583	
18	-2.7	-2.7	0.01042093481	
19	-2.6	-2.6	0.01358296923	
20	-2.5	-2.5	0.01752830049	
21	-2.4	-2.4	0.02239453029	

Step 5: Visualizing the Data using a Scatter Chart

The final step involves plotting the calculated X and Y coordinates to visualize the normal distribution. Since we are plotting continuous mathematical coordinates rather than discrete categories, the ideal visualization tool is the **Scatter Chart**.

Follow these steps to generate the visualization:

Highlight the entire range containing your X-values and Y-values (in our example, **B5:C85**).

Navigate to the top menu and click **Insert**.

Select **Chart** from the dropdown menu.


Google Sheets may default to a Line Chart; however, for precise mathematical plots, we must ensure the Chart Type is set to **Scatter Chart** (or a Scatter Chart with smoothed lines). This will connect the dense collection of data points, producing the unmistakable **bell curve** shape.



-2.3	0.02832703774					
-2.2	0.03547459285					
-2.1	0.04398359598					
-2	0.05399096651					
-1.9	0.06561581477					
-1.8	0.0789501583					
-1.7	0.09404907738					

ates and au
if

if



To enhance readability and professional appearance, utilize the Chart Editor to refine the visualization. Best practices include modifying the chart title, ensuring axis labels accurately describe the data (e.g., "Probability Density" for the Y-axis and "Data Value" for the X-axis), and selecting appropriate colors and line thicknesses.

Conclusion and Next Steps

By leveraging the statistical power of the NORMDIST function and the flexibility of the Scatter Chart type in Google Sheets, you can efficiently generate accurate and dynamic visualizations of any **bell curve** distribution. This method is indispensable for data analysis, grading curves, and probability modeling.

For those looking to deepen their understanding of statistical distributions and related concepts, we recommend the following resources:

[An Introduction to the Normal Distribution](#)

[Normal CDF Calculator](#)