

How to Calculate Line of Best Fit on a TI-84 Calculator: A Step-by-Step Guide

Authored by
stats writer

December 5, 2025

RECOMMENDED CITATION

stats writer (2025). *How to Calculate Line of Best Fit on a TI-84 Calculator: A Step-by-Step Guide*. PSYCHOLOGICAL SCALES. Retrieved from <https://scales.arabpsychology.com/?p=105775>

The process of determining the line of best fit, also known as linear regression, is a fundamental technique in statistics used to model the relationship between two variables. Utilizing a powerful graphing calculator like the TI-84 calculator streamlines this complex calculation, providing immediate graphical and numerical results. Before beginning, it is essential to understand that finding this specific line involves minimizing the sum of the squared residuals, ensuring the line truly represents the central trend of the data points.

Once the requisite data has been meticulously entered into the calculator's lists, the user navigates through the statistical calculation menus. Specifically, selecting the "Stat" menu, moving to the "Calc" submenu, and choosing the appropriate linear regression model--either "LinReg(a+bx)" or "LinReg(ax+b)"--initiates the calculation. The choice between these two forms usually depends on the convention preferred in a specific curriculum, but mathematically, they yield identical results for the relationship. Upon execution, the calculator outputs the equation coefficients, the correlation coefficient (r), and the coefficient of determination (r-squared), providing a comprehensive statistical summary of the relationship under scrutiny.

A **line of best fit**, formally known as the least-squares regression line, represents the best linear model that describes the relationship between an independent variable (x) and a dependent variable (y) within a given dataset. This line is crucial for predicting future values and quantifying the strength and direction of the linear association between the variables.

This tutorial provides a meticulous, step-by-step example detailing how to utilize the advanced functions of the TI-84 graphing calculator to calculate the line of best fit for a sample dataset. For demonstration purposes, we will use the following paired data:

x	y
2	8
4	9
5	12
6	14
8	13
11	19

Understanding the Theory of Linear Regression

Before diving into the mechanics of the calculator, grasping the underlying principles of linear regression is vital. Linear regression seeks to model the relationship between two variables by fitting a straight line to observed data. The general form of the line is often expressed as $y = ax +$

$y = a + bx$. The coefficient 'a' represents the slope of the line, indicating the change in the dependent variable (y) for every one-unit change in the independent variable (x), while 'b' represents the y-intercept, the value of y when x is zero.

The calculation performed by the TI-84 is known as the Ordinary Least Squares (OLS) method. This method determines the unique line that minimizes the sum of the squared vertical distances (residuals) from each data point to the line itself. This minimization process ensures that the resulting line is optimally positioned to represent the overall trend of the data, thereby minimizing prediction errors. When interpreting the final results, always remember that correlation does not necessarily imply causation, even when a strong line of best fit is established.

Furthermore, the calculator provides two crucial metrics for assessing the quality of the fit: the correlation coefficient (r) and the coefficient of determination (r^2). The value of 'r' ranges from -1 to 1, indicating the strength and direction of the linear relationship. A value close to 1 indicates a strong positive linear relationship, while a value close to -1 indicates a strong negative linear relationship. A value near zero suggests little to no linear relationship. The r^2 value, conversely, represents the proportion of the variance in the dependent variable that is predictable from the independent variable, offering a measure of how well the regression predictions approximate the real data points. Understanding these metrics is essential for accurate statistical analysis.

Step 1: Preparing the Calculator and Enabling Diagnostics

To ensure that your TI-84 provides the complete statistical output, including the correlation coefficient (r) and the coefficient of determination (r^2), it is often necessary to activate the Diagnostic feature. While modern TI-84 models often have this setting enabled by default, older versions or recently reset calculators require manual activation. Failure to perform this check may result in the calculator only displaying the 'a' and 'b' coefficients without the crucial goodness-of-fit indicators.

To enable Diagnostics, first press the 2nd key, followed by the 0 key (which accesses the CATALOG menu). The CATALOG contains every function available on the calculator, listed alphabetically. Scroll down extensively until you locate the function labeled DiagnosticOn. Press ENTER to select this function, and then press ENTER again to execute the command. The screen will momentarily display "Done," confirming that the calculator is now configured to show 'r' and ' r^2 ' during regression calculations.

It is good practice to clear any previous data from the lists before starting a new regression analysis to prevent confusion or accidental inclusion of irrelevant data points. To clear the lists, press STAT, navigate to the EDIT menu, and select option 4, ClrList. After selecting this option, specify the lists you wish to clear, typically L1 and L2 (accessed by pressing 2nd and 1, followed by a comma, and then 2nd and 2). Executing this command ensures a clean slate for the current


dataset, improving the reliability and integrity of the subsequent regression analysis.

Step 2: Entering and Managing the Dataset

The initial and most critical phase of statistical analysis on the TI-84 is accurate data entry. Any errors made during this step will render the resulting line of best fit and associated statistics invalid. This process requires organizing the independent variable data (x-values) into one list and the dependent variable data (y-values) into a corresponding list, ensuring that the paired values align horizontally.

To begin data entry, press the dedicated STAT button, which accesses all statistical functionalities. From the subsequent menu, select option 1, which is EDIT. This action opens the list editor, typically displaying lists L1, L2, L3, and so on. We will use L1 for the independent x-values and L2 for the dependent y-values. Carefully input each x-value into column L1, pressing ENTER after each entry to move to the next row. Once L1 is complete, use the right arrow key to move the cursor to the top of column L2.

Proceed to enter the corresponding y-values into column L2, maintaining the precise order of the pairings. It is imperative that both lists, L1 and L2, contain exactly the same number of data points; a mismatch will cause a "DIMENSION MISMATCH" error when attempting the regression calculation. After entering all values, take a moment to double-check the entered data against the source table to confirm accuracy before proceeding to the calculation phase. For the example dataset provided earlier, your screen should appear as follows:




Step 3: Executing the Linear Regression Calculation (LinReg)

With the data correctly stored in L1 and L2, the next step is to instruct the TI-84 calculator to

perform the linear regression analysis. This calculation is accessed through the statistical test menu, which houses all available regression and analysis functions. This is where the magic of the least-squares method is applied to your data.

Press the STAT button once more, but instead of selecting EDIT, use the right arrow key to scroll over to the CALC menu. The CALC menu contains various regression models (linear, quadratic, exponential, power, etc.). Scroll down through the options until you locate the linear regression function. You will typically see two options: LinReg(ax+b) (Option 4) and LinReg(a+bx) (Option 8). Select LinReg(ax+b), which is the standard notation in many educational settings, and press ENTER. The display should show the function on the home screen.

Modern TI-84 models (such as the Plus CE) will present a setup screen requiring input for Xlist, Ylist, FreqList, and Store RegEQ. For older models, you must manually input the lists and the location to store the equation, separated by commas. Specify L1 for the Xlist and L2 for the Ylist. Crucially, the "Store RegEQ" function allows the calculated regression equation to be automatically saved into the Y= function editor, which is necessary for plotting the line later. To store the equation, position the cursor next to "Store RegEQ," and press VARS, scroll right to Y-VARS, select Option 1: Function, and then select Y1. Your setup should look like the following image, ensuring the calculator knows where to find the data and where to store the result:



Finally, scroll down to Calculate and press ENTER. The TI-84 will instantly process the data points using the OLS method and display the detailed results screen.


Step 4: Interpreting the Regression Output

The results screen generated by the calculator contains all the statistical information required to define and assess the quality of the line of best fit. This output is presented in the form of equation parameters and crucial diagnostic statistics, which must be correctly interpreted to draw valid conclusions about the dataset.

The calculator first defines the form of the equation: $y = ax + b$. It then provides the numerical values for the coefficient a (the slope) and the coefficient b (the y-intercept). For our example

data, the output shows: \$a\$ approx 1.14\$ and \$b\$ approx 5.493\$. Therefore, the calculated line of best fit is: $y = 1.14x + 5.493$. This means that for every one-unit increase in the independent variable (x), the dependent variable (y) increases by approximately 1.14 units. The y -intercept of 5.493 represents the predicted value of y when x is zero.

The screen also provides the two critical diagnostic values (assuming diagnostics were enabled in Step 1). The correlation coefficient (r) is typically displayed next, indicating the strength and direction of the linear relationship. For our example, \$r\$ approx 0.9928\$. Because this value is extremely close to +1, it signifies a very strong positive linear correlation between the x and y variables. This high 'r' value suggests that the linear model is highly appropriate for this data. The second crucial value is the coefficient of determination (r^2), which represents the proportion of variability in Y explained by the linear relationship with X . Our example shows \$r^2\$ approx 0.9857\$. This indicates that approximately 98.57% of the variation in y can be explained by the variation in x using this linear model, suggesting an exceptional fit. The full output screen for the calculation is shown below:



The calculated line of best fit is: $y = 5.493 + 1.14x$ (Note: The TI-84 output typically lists 'a' first, followed by 'b', matching the $y = ax + b$ format. When writing the equation, we maintain this structure or use the equivalent $y = b + ax$ form as shown in the original content.)


Step 5: Plotting the Data and the Regression Line

While the numerical output is essential, visualizing the data and the calculated line of best fit provides an indispensable check on the analysis. Plotting the data confirms that the linear model visually matches the scattered points, ensuring there are no obvious non-linear patterns or influential outliers that might skew the results. Since we stored the regression equation in $Y1$ during Step 3, the calculator is already prepared to graph the line.

First, ensure the calculator is set up to display the scatterplot. Press the 2nd key, followed by Y= (STAT PLOT menu). Select Plot 1 and ensure it is set to ON, with Type set to Scatterplot (the first option), Xlist set to L1, and Ylist set to L2. This configuration prepares the calculator to draw the individual data points.

Next, use the specialized statistical zoom function to automatically adjust the viewing window to optimally fit all data points. Press the ZOOM key, which offers various graphing window options. Scroll down the menu until you find option 9, labeled ZoomStat. Press ENTER. This command scales the x and y axes precisely to encompass the minimum and maximum values of your L1 and L2 data, simultaneously plotting the data points and the regression line stored in Y1.

The resulting display will show the scatterplot with the straight line of best fit superimposed directly over the data points. For our example, the strong correlation (r close to 1) means the points should cluster very closely around the plotted line, visually confirming the high coefficient of determination (r^2) found in the previous step. The resulting graphical output should look like the following:



Conclusion: Leveraging Statistical Insights

The ability to quickly and accurately calculate the line of best fit using the TI-84 calculator is an invaluable skill for students and professionals alike. This procedure not only yields the predictive equation but also provides critical metrics, such as the correlation coefficient (r), that quantify the reliability of the model. By following these structured steps--entering data, executing the LinReg command, and interpreting the output--you can effectively analyze linear relationships within any bivariate dataset.

It is important to remember that the validity of the linear model depends heavily on the assumption that the relationship between the variables is indeed linear. Always examine the scatterplot; if the

data points suggest a curved trend (e.g., quadratic or exponential), a different regression model should be selected from the CALC menu for a more accurate fit. Mastery of the TI-84's statistical functions allows for rapid exploration of various models, leading to robust statistical conclusions.

For more complex analyses or when dealing with multiple explanatory variables (multiple regression), specialized statistical software is necessary. However, for introductory statistics and two-variable analysis, the TI-84 remains an indispensable tool for generating reliable least-squares regression lines and their associated measures of fit and correlation.

ARABPSYCHOLOGY.COM