

What is the process for finding the area to the left of a given z-score, and what are some examples of how it is used?

Authored by
stats writer

May 5, 2024

RECOMMENDED CITATION

stats writer (2024). *What is the process for finding the area to the left of a given z-score, and what are some examples of how it is used?*. PSYCHOLOGICAL SCALES. Retrieved from <https://scales.arabpsychology.com/?p=142998>

The process for finding the area to the left of a given z-score involves using the standard normal distribution table or a statistical software to determine the probability of obtaining a value less than the given z-score. This area is also known as the cumulative probability or the p-value.

This process is commonly used in statistics to analyze and interpret data. For example, in hypothesis testing, the area to the left of a certain z-score is used to determine the significance of a test statistic. A smaller area indicates a stronger evidence against the null hypothesis, while a larger area suggests weaker evidence.

In addition, the area to the left of a z-score is also used in calculating confidence intervals, where it represents the probability of obtaining a value within a certain range.

Overall, the process of finding the area to the left of a given z-score is a fundamental step in many statistical analyses and helps to provide insights into the likelihood of a particular outcome occurring.

Find Area to the Left of Z-Score (With Examples)

In statistics, a z-score tells us how many standard deviations away a given value lies from a population mean.

We use the following formula to calculate a z-score for a given value:

$$z = (x - \mu) / \sigma$$

where:

x: Individual data value
 μ : Mean of population
 σ : Standard deviation of population

To find the area under a normal distribution that lies to

the left of a given z-score, we can use one of two methods:

1. Use the .

2. Use the .

The following examples show how to use each of these methods in practice.

Example 1: Area to the Left of Negative Z-Score

The weight of a certain species of turtles is normally distributed with mean $\mu = 300$ pounds and standard deviation $\sigma = 15$ pounds. Approximately what percentage of turtles weigh less than 284 pounds?

The z-score for a weight of 284 pounds would be calculated as $z = (284 - 300) / 15 = -1.07$

We can use one of two methods to find the area to the left of this z-score:

Method 1: Use z table.

To find the area to the left of the z-score, we can simply look up the value -1.07 in the :

z	0	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
-3.0	0.0013	0.0013	0.0013	0.0012	0.0012	0.0011	0.0011	0.0011	0.0010	0.0010
-2.9	0.0019	0.0018	0.0018	0.0017	0.0016	0.0016	0.0015	0.0015	0.0014	0.0014
-2.8	0.0026	0.0025	0.0024	0.0023	0.0023	0.0022	0.0021	0.0021	0.0020	0.0019
-2.7	0.0035	0.0034	0.0033	0.0032	0.0031	0.0030	0.0029	0.0028	0.0027	0.0026
-2.6	0.0047	0.0045	0.0044	0.0043	0.0041	0.0040	0.0039	0.0038	0.0037	0.0036
-2.5	0.0062	0.0060	0.0059	0.0057	0.0055	0.0054	0.0052	0.0051	0.0049	0.0048
-2.4	0.0082	0.0080	0.0078	0.0075	0.0073	0.0071	0.0069	0.0068	0.0066	0.0064
-2.3	0.0107	0.0104	0.0102	0.0099	0.0096	0.0094	0.0091	0.0089	0.0087	0.0084
-2.2	0.0139	0.0136	0.0132	0.0129	0.0125	0.0122	0.0119	0.0116	0.0113	0.0110
-2.1	0.0179	0.0174	0.0170	0.0166	0.0162	0.0158	0.0154	0.0150	0.0146	0.0143
-2.0	0.0228	0.0222	0.0217	0.0212	0.0207	0.0202	0.0197	0.0192	0.0188	0.0183
-1.9	0.0287	0.0281	0.0274	0.0268	0.0262	0.0256	0.0250	0.0244	0.0239	0.0233
-1.8	0.0359	0.0351	0.0344	0.0336	0.0329	0.0322	0.0314	0.0307	0.0301	0.0294
-1.7	0.0446	0.0436	0.0427	0.0418	0.0409	0.0401	0.0392	0.0384	0.0375	0.0367
-1.6	0.0548	0.0537	0.0526	0.0516	0.0505	0.0495	0.0485	0.0475	0.0465	0.0455
-1.5	0.0668	0.0655	0.0643	0.0630	0.0618	0.0606	0.0594	0.0582	0.0571	0.0559
-1.4	0.0808	0.0793	0.0778	0.0764	0.0749	0.0735	0.0721	0.0708	0.0694	0.0681
-1.3	0.0968	0.0951	0.0934	0.0918	0.0901	0.0885	0.0869	0.0853	0.0838	0.0823
-1.2	0.1151	0.1131	0.1112	0.1093	0.1075	0.1056	0.1038	0.1020	0.1003	0.0985
-1.1	0.1357	0.1335	0.1314	0.1292	0.1271	0.1251	0.1230	0.1210	0.1190	0.1170
-1.0	0.1587	0.1562	0.1539	0.1515	0.1492	0.1469	0.1446	0.1423	0.1401	0.1379
-0.9	0.1841	0.1814	0.1788	0.1762	0.1736	0.1711	0.1685	0.1660	0.1635	0.1611
-0.8	0.2119	0.2090	0.2061	0.2033	0.2005	0.1977	0.1949	0.1922	0.1894	0.1867
-0.7	0.2420	0.2389	0.2358	0.2327	0.2296	0.2266	0.2236	0.2206	0.2177	0.2148
-0.6	0.2743	0.2709	0.2676	0.2643	0.2611	0.2578	0.2546	0.2514	0.2483	0.2451

The area to the left of $z = -1.07$ is 0.1423.

Method 2: Use Area to the Left of Z-Score Calculator

We can also use the to find that the area to the left of $z = -1.07$ is 0.1423.

Area To The Left of Z-Score Calculator

This calculator finds the area to the left of a certain **z-score** in the normal distribution.

Simply enter the z-score below and then click the "Calculate" button.

Z-Score

CALCULATE

Area to the Left of Z-Score: 0.14231

Example 2: Area to the Left of Positive Z-Score

The scores on a certain exam are normally distributed with mean $\mu = 85$ and standard deviation $\sigma = 8$. Approximately what percentage of students score less than 87 on the exam?

The z-score for an exam score of 87 would be calculated as $z = (87 - 85) / 8 = 0.25$

We can use one of two methods to find the area to the left of this z-score:

Method 1: Use z table.

To find the area to the left of the z-score, we can simply look up the value 0.25 in the :

z	0	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545

The area to the left of $z = 0.25$ is 0.5987. Applied to our scenario, this means approximately 59.87% of students score less than 87 on this exam.

Method 2: Use Area to the Left of Z-Score Calculator

We can also use the to find that the area to the left of $z = 0.25$ is 0.5987.

Area To The Left of Z-Score Calculator

This calculator finds the area to the left of a certain **z-score** in the normal distribution.

Simply enter the z-score below and then click the "Calculate" button.

Z-Score

CALCULATE

Area to the Left of Z-Score: 0.59871

The following tutorials provide additional information on how to work with z-scores: