

DONG QUAI

Authored by
mohammad looti

October 30, 2025

RECOMMENDED CITATION

mohammad looti (2025). *DONG QUAI*. PSYCHOLOGICAL SCALES. Retrieved from <https://scales.arabpsychology.com/?p=64542>

Dong Quai

Primary Disciplinary Field(s): Pharmacology, Herbal Medicine, Traditional Chinese Medicine (TCM)

1. Core Definition and Nomenclature

Dong Quai (simplified Chinese: 当归; pinyin: dāngguī) is an extensively studied and utilized **herbal agent** derived from the dried root of the plant species *Angelica sinensis* (Oliv.) Diels, a member of the Apiaceae family. This botanical substance is highly esteemed within the materia medica of Traditional Chinese Medicine (TCM), where it is often referred to as the "female ginseng" due to its widespread application in treating various gynecological conditions. The term **Dong Quai** itself translates roughly to "state of return" or "ought to return," metaphorically referencing the belief that the herb restores health and balance, particularly regarding reproductive health and blood flow. Its significance extends beyond gynecological applications, often being included in complex herbal formulas aimed at tonifying the blood, invigorating circulation, and alleviating pain.

The preparation of Dong Quai involves harvesting the roots, typically during late autumn, followed by drying and sometimes processing with wine (alcohol) to enhance specific therapeutic properties according to TCM principles. While the primary source is *Angelica sinensis*, related species such as *Angelica acutiloba* (Japanese Angelica) and *Angelica gigas* (Korean Angelica) are used in regional variations of traditional medicine, though **Dong Quai** specifically refers to the Chinese variety. The standardization of the herbal product is critical for modern scientific study, often relying on quantifying key bioactive compounds such as ferulic acid and ligustilide to ensure consistency in pharmacological efficacy across different batches and preparations.

In academic and clinical contexts outside of TCM, Dong Quai is classified as a botanical dietary supplement. Its increasing global prominence has driven substantial research into its chemical composition and mechanisms of action, attempting to bridge the gap between empirical traditional use and evidence-based pharmacology. This research is particularly focused on understanding the complex interplay of its numerous constituents and their collective impact on the endocrine, hematopoietic, and cardiovascular systems. The inherent complexity of herbal medicines, where the therapeutic effect is often attributed to synergy rather than a single compound, makes the scientific validation of **Dong Quai's** efficacy both challenging and necessary.

2. Botanical Origin and Geographical Distribution

The plant from which Dong Quai is derived, *Angelica sinensis*, is a perennial herbaceous plant indigenous to the high altitude, cool, and damp mountainous regions of East Asia. Specifically, its primary native geographical distribution encompasses the provinces of Gansu, Shaanxi, Hubei,

Sichuan, and Yunnan in **China**, as well as parts of **Korea** and **Japan**. Due to high demand in herbal medicine markets, *Angelica sinensis* is now extensively cultivated across these regions, though the quality of the root is often thought to be highest when harvested from its original native environments, particularly in the mountainous regions of western China.

Botanically, *Angelica sinensis* grows to heights of up to two meters, characterized by large tripinnate leaves and compound umbels of small white flowers. The root, which constitutes the medicinal part, is thick, fleshy, and highly aromatic, possessing a distinct sweet, pungent, and slightly bitter flavor profile. The quality assessment in TCM often considers factors such as root size, color, internal structure (being fatty and smooth), and fragrance, which are believed to correlate with its clinical effectiveness. Proper growth conditions, including adequate moisture and specific soil nutrients, are crucial for the development of roots rich in the desired volatile oils and non-volatile compounds responsible for its therapeutic activities.

The geographical specificity of cultivation influences the chemical fingerprint of the harvested root. Variations in climate, altitude, and cultivation techniques can lead to significant differences in the concentration of key bioactive compounds, such as ligustilide, ferulic acid, and various polysaccharides. These differences sometimes necessitate regional adjustments in dosage or preparation methods when utilized in traditional pharmacopoeias. The global trade of **Dong Quai** requires careful documentation and quality control to ensure that products sourced internationally maintain the phytochemical profile necessary for the expected pharmacological outcomes, highlighting the importance of tracing the botanical origin back to reliable cultivation sites.

3. Traditional Chinese Medicine (TCM) Applications

Within the framework of TCM, Dong Quai is fundamentally categorized as a herb that **nourishes and harmonizes the blood** (B? Xuè), regulates the menses (Tiáo J?ng), and disperses cold to stop pain (Q? Hán Zh? Tòng). It is considered warm in nature and sweet and pungent in taste, acting predominantly on the Liver, Heart, and Spleen channels. Its primary therapeutic role revolves around addressing conditions characterized by "Blood Deficiency" (Xuè X?) or "Blood Stasis" (Xuè Y?), which often manifest with symptoms of pallor, fatigue, dry skin, and irregular or painful menstruation.

The most traditional and widespread use of **Dong Quai**, confirmed by the source content, is in addressing **menstruation problems**. This includes the treatment of dysmenorrhea (painful periods), amenorrhea (absence of periods), and premenstrual syndrome (PMS). TCM practitioners often prescribe Dong Quai in combinations, such as the famous formula Si Wu Tang (Four-Substance Decoction), where it synergizes with other herbs to powerfully replenish blood while ensuring adequate circulation. Its reputation as a powerful regulator of female reproductive health has cemented its status, historically being the most frequently prescribed herb for managing

climacteric symptoms and postpartum recovery due to its perceived ability to rebuild blood lost during childbirth.

Beyond gynecological health, **Dong Quai** is also employed for its analgesic (pain-relieving) properties, particularly for chronic pain associated with poor circulation, abdominal stagnation, and traumatic injuries. Furthermore, its blood-tonifying function makes it useful in treating anemia, chronic fatigue, and general weakness. TCM theory posits that while it replenishes blood, its volatile oil content ensures that the blood moves smoothly, preventing stagnation. This dual action-tonifying without causing stasis--is highly valued and distinguishes it from some other tonic herbs. Its use also extends to treating constipation, especially in the elderly or those with dryness, due to its mild laxative effect and ability to moisten the intestines.

4. Phytochemistry and Active Components

The complex therapeutic profile of Dong Quai is attributable to its diverse and extensive phytochemical composition, which includes volatile oils, organic acids, phthalides, polysaccharides, and coumarins. The volatile fraction, which contributes significantly to the root's characteristic aroma, is rich in **phthalides**, most notably ligustilide, butylphthalide, and sedanenolide. Ligustilide, in particular, is considered a key quality marker and is responsible for many of the antispasmodic and smooth muscle relaxant properties that contribute to its efficacy in treating menstrual cramps and circulatory issues.

Another critical non-volatile component is **ferulic acid**, an organic acid widely recognized for its strong antioxidant and anti-inflammatory effects. Ferulic acid plays a major role in the herb's cardiovascular benefits and is often standardized in commercial extracts. Additionally, the root contains various coumarins and furanocoumarins, which are compounds known for their potential influence on blood coagulation. It is the presence of these coumarin derivatives that gives rise to the herb's known **blood-thinning properties**, a critical factor in its safety profile and potential for drug interactions.

In addition to these low-molecular-weight compounds, **Dong Quai** contains high-molecular-weight polysaccharides. These polysaccharides have been the focus of modern research regarding the herb's immunomodulatory effects, showing potential benefits in stimulating hematopoietic recovery and regulating immune responses. The synergy between the volatile oils (promoting movement and reducing spasm), ferulic acid (providing protection), and the polysaccharides (modulating immunity and blood production) exemplifies why the whole herb, rather than isolated compounds, is often preferred in traditional contexts. Understanding this complex chemical matrix is essential for fully evaluating the herb's efficacy and ensuring safety in clinical application.

5. Pharmacological Mechanisms of Action

The pharmacological actions of **Dong Quai** are highly diverse, reflecting its complex chemical makeup and its broad traditional applications. One of the best-studied mechanisms relates to its effect on smooth musculature. Ligustilide and other phthalides exhibit potent **antispasmodic activity**, particularly on the uterine muscles, which is thought to be the primary mechanism by which Dong Quai alleviates dysmenorrhea. By relaxing the smooth muscles of the uterus, it can reduce painful contractions and improve localized blood flow, addressing both the pain and the stasis components of menstrual distress.

Furthermore, Dong Quai possesses significant effects on the **hematopoietic system** and circulation. Studies indicate that it can promote hematopoiesis (the formation of blood cellular components), supporting its traditional use in treating anemia and blood deficiency. Simultaneously, its compounds, particularly certain coumarins, demonstrate measurable **anticoagulant activity**. This anticoagulant effect is crucial to understand, as the source material warns that the herb "contains blood thinning properties." This action helps to break up blood stasis (Xuè Y?) and improve overall microcirculation, which is beneficial for pain relief and tissue recovery, but carries risks, as detailed below.

Beyond the physical effects on blood and muscles, the components of **Dong Quai** also display neuroprotective, antioxidant, and anti-inflammatory properties. Ferulic acid is a strong radical scavenger, mitigating oxidative stress associated with various chronic diseases. The herb has also been shown to influence hormonal balance, though the mechanism is not estrogenic in the classical sense. While earlier research suggested phytoestrogenic effects, subsequent comprehensive studies indicate that **Dong Quai** does not typically bind to estrogen receptors but may modulate endogenous hormone levels or mimic certain estrogenic effects through non-receptor mediated pathways, supporting its traditional use in menopausal symptom management.

6. Safety, Drug Interactions, and Contraindications

Despite its long history of safe use in traditional practice, **Dong Quai** is not without significant safety considerations, particularly regarding its interactions with pharmaceutical drugs and its effect on coagulation. The most critical caution, explicitly noted in the source material, is related to its **blood-thinning properties**. Due to the presence of coumarins, Dong Quai may potentiate the effects of anticoagulant and antiplatelet medications, such as Warfarin, heparin, and aspirin. Concurrent use can significantly increase the risk of bleeding or hemorrhage. Therefore, the advisory to only take Dong Quai after consulting a doctor is paramount, especially for individuals already on anticoagulant therapy or those scheduled for surgery.

Another significant safety concern involves the photosensitivity induced by certain furanocoumarins

present in the root. These compounds can increase the skin's sensitivity to ultraviolet light, potentially leading to photodermatitis or sunburn, particularly in fair-skinned individuals or during periods of prolonged sun exposure. Users taking **Dong Quai** should be advised to take protective measures against the sun. Furthermore, while generally well-tolerated, side effects such as gastrointestinal upset, bloating, and allergic reactions (rarely) have been reported.

Contraindications for the use of **Dong Quai** are generally strict for certain populations. It is contraindicated in pregnant women due to potential uterine stimulating effects, particularly at high doses, although traditional use often included formulations for post-partum recovery. It should also be avoided by individuals with bleeding disorders, heavy menstrual bleeding, or those with hormone-sensitive cancers (due to the previously mentioned, albeit weak or non-classical, hormonal modulating potential). Due to the complex nature of herbal medicine interaction, high-quality clinical guidance is essential, reinforcing the necessity that "A person should only take dong quai if they let their doctor know."

Further Reading

[Wikipedia Entry: Angelica sinensis \(Dong Quai\)](#)

[National Institutes of Health \(NIH\): Review of Angelica sinensis for dysmenorrhea](#)

[Wikipedia Entry: Traditional Chinese Medicine \(TCM\)](#)

[WebMD: Dong Quai Safety and Dosage Information](#)