

# FIGHT-FLIGHT REACTION

Authored by  
**mohammad looti**

October 11, 2025

## RECOMMENDED CITATION

mohammad looti (2025). *FIGHT-FLIGHT REACTION*. PSYCHOLOGICAL SCALES.  
Retrieved from <https://scales.arabpsychology.com/?p=43343>

## Fight-Flight Reaction

**Primary Disciplinary Field(s):** Psychology, Physiology, Neurobiology

### 1. Core Definition

The **fight-flight reaction**, formally known as the acute stress response or hyperarousal, is an automatic, non-conscious physiological mechanism that occurs in response to an event perceived as physically or psychologically harmful, threatening, or terrifying. This survival mechanism is designed to prepare the organism to either vigorously combat the perceived threat (fight) or rapidly flee from it (flight). It is often referred to as the emergency reaction or emergency syndrome, emphasizing its crucial role in immediate survival.

This reaction is fundamentally mediated by the sympathetic nervous system, a division of the autonomic nervous system. When a threat is detected--whether real (e.g., a physical attack) or perceived (e.g., intense public speaking anxiety)--the hypothalamus initiates a cascade of hormonal releases. These hormones, primarily cortisol and adrenaline (epinephrine), induce a rapid systemic shift, diverting energy resources away from long-term maintenance processes (like digestion and immunity) toward immediate muscular and neurological readiness.

The speed and efficiency of the fight-flight reaction are paramount. It bypasses rational, reflective thought processes controlled by the prefrontal cortex, instead relying on the rapid processing capabilities of the amygdala. This evolutionary necessity allows for instantaneous action when hesitation could result in serious injury or death, ensuring that the organism prioritizes survival over complex cognitive assessment during moments of acute danger.

### 2. Etymology and Historical Development

While humans have always experienced intense, primal responses to danger, the scientific understanding and naming of this specific mechanism are largely attributed to American physiologist **Walter Bradford Cannon**. Working in the early 20th century, Cannon established the crucial relationship between emotional stress and physiological responses, formalizing his observations in his 1915 work, *Bodily Changes in Pain, Hunger, Fear and Rage*.

Cannon's research focused heavily on the role of the sympathetic nervous system and the rapid release of catecholamines, demonstrating how these biological factors prepared the body for energy expenditure. His work provided the foundational framework for modern stress theory, emphasizing the body's innate capacity for maintaining internal stability--a concept he termed homeostasis. He argued that the fight-flight response was the necessary mechanism to restore this internal balance following external disruption.

In subsequent decades, the model was refined. Notably, researchers recognized that the reaction to danger often involved a third, equally adaptive response: **freeze**. The inclusion of the freeze response acknowledges that in certain situations, particularly when escape or confrontation is impossible, tonic immobility (playing dead or remaining perfectly still) may be the most advantageous survival strategy, especially in predatory encounters where movement triggers attack. Thus, the concept is now frequently discussed as the **fight-flight-freeze response**.

### 3. Key Concepts and Components

The fight-flight reaction involves coordinated activation across multiple physiological systems, controlled by the hypothalamic-pituitary-adrenal (HPA) axis and the autonomic nervous system. The immediate preparation for emergency action relies on several integrated components:

**Sympathetic Nervous System Activation:** The initial trigger involves the immediate activation of the sympathetic nervous system, causing the adrenal glands to release adrenaline (epinephrine) and noradrenaline (norepinephrine). These hormones travel rapidly through the bloodstream, initiating widespread physical changes.

**Cardiovascular and Respiratory Changes:** Heart rate increases dramatically (tachycardia) and blood pressure rises, accelerating the transport of oxygen and glucose to the major muscle groups. Respiration quickens (hyperventilation), ensuring maximum oxygen uptake to fuel the anticipated intense physical exertion required for fighting or fleeing.

**Musculoskeletal Mobilization:** Blood flow is strategically redirected from non-essential areas (such as the skin, digestive tract, and reproductive organs) to the skeletal muscles, preparing them for immediate, powerful action. This shunting of blood also causes paleness and peripheral coldness.

**Sensory and Cognitive Alterations:** Pupils dilate (mydriasis) to allow maximum light intake, enhancing vision for identifying threats. Pain perception temporarily diminishes (stress-induced analgesia) to allow the organism to continue fighting or running despite injury. Cognitive focus narrows intensely on the threat, sometimes leading to "tunnel vision," while peripheral awareness is suppressed.

**Metabolic Shifts:** The liver converts stored glycogen into glucose, which is released into the bloodstream, providing a surge of metabolic fuel necessary for the high-energy demands of the emergency response.

### 4. Physiological Cascade

The physiological process follows a predictable cascade once a threat is registered. First, sensory information is processed instantly by the thalamus, which routes the input simultaneously to the slower, analytical cortex and the faster, emotional amygdala. The amygdala, recognizing danger based on previous memory or instinct, triggers the hypothalamus.

The hypothalamus acts as the command center, initiating two main pathways: the rapid sympathetic-adreno-medullary (SAM) axis and the slower, but more sustained, HPA axis. The SAM axis is responsible for the adrenaline rush and the immediate physical changes--this is the "fight-flight" part. The HPA axis involves the release of corticotropin-releasing hormone (CRH), which stimulates the pituitary gland to release adrenocorticotropic hormone (ACTH). ACTH travels to the adrenal cortex, prompting the release of **cortisol**.

Cortisol is the primary stress hormone responsible for maintaining the heightened state of arousal over a longer duration, ensuring sufficient energy reserves are available. Once the threat passes, the parasympathetic nervous system (the "rest and digest" system) engages, releasing neurotransmitters like acetylcholine to slow the heart rate, normalize breathing, and gradually return the body to a state of homeostasis.

## 5. Significance and Impact

The fight-flight reaction holds immense evolutionary significance. For early humans and non-human species, its functionality was absolute: it provided the crucial edge necessary for surviving predation or hostile encounters. Organisms lacking this rapid response mechanism would quickly be eliminated from the gene pool.

In the modern context, however, the reaction's adaptive value is complicated. While essential for responding to acute physical emergencies (e.g., stopping short of an oncoming car), the reaction is frequently triggered by chronic, non-lethal psychological stressors--such as workplace deadlines, financial anxiety, or relationship conflicts. Since the body is constantly prepared for physical combat or escape but cannot utilize that energy effectively (one cannot physically fight a bad credit score), the sustained activation of the HPA axis can become detrimental.

Chronic activation of the fight-flight response leads to persistently high levels of cortisol and adrenaline, contributing significantly to modern health issues. These include anxiety disorders, hypertension, cardiovascular disease, immune system suppression, gastrointestinal problems, and chronic fatigue. Understanding the mechanism is therefore critical in clinical psychology and medicine for treating stress-related illnesses.

## 6. Debates and Extended Models

While the fight-flight model remains the cornerstone of stress physiology, it has been subject to expansion and refinement over time to account for complex behavioral responses.

**The Tend-and-Befriend Response:** Sociological and psychological research, particularly led by Shelley Taylor and colleagues, introduced the tend-and-befriend model, primarily observed in females. This model suggests that, particularly under stress, females may exhibit behaviors aimed

at protecting offspring (tending) and seeking social support or forming alliances (befriending) rather than immediate flight or fight. This response is theorized to be modulated by oxytocin, which counteracts the immediate effects of cortisol and adrenaline.

**The Inclusion of Freeze:** The addition of the "freeze" response is now widely accepted. This response, often linked to dissociation or tonic immobility, is particularly relevant in the study of trauma and post-traumatic stress disorder (PTSD). During freeze, heart rate and respiration may slow dramatically, and the individual may appear disconnected or unresponsive, often an adaptive response when overwhelmed by a perceived unavoidable threat.

**The Faint Response:** Some researchers also include the "faint" response, a vasovagal syncope, which causes a sudden drop in heart rate and blood pressure. While less common, this response may serve an evolutionary purpose by reducing blood loss if injured, though it is often considered a maladaptive overreaction in non-physical threat scenarios.

## 7. Further Reading

Walter Cannon (Physiologist)

Sympathetic Nervous System

Adrenaline (Epinephrine)

Tend and Befriend Theory