

# PERIRHINAL CORTEX

Authored by  
**mohammad looti**

October 27, 2025

## RECOMMENDED CITATION

mohammad looti (2025). *PERIRHINAL CORTEX*. PSYCHOLOGICAL SCALES. Retrieved from <https://scales.arabpsychology.com/?p=60739>

## PERIRHINAL CORTEX

**Primary Disciplinary Field(s):** Neuroscience, Cognitive Psychology, Neuroanatomy

### 1. Core Definition

The **Perirhinal Cortex** (PRC), corresponding primarily to Brodmann areas 35 and 36, is an essential structure located within the Medial Temporal Lobe (MTL), situated adjacent to the hippocampus and the entorhinal cortex. Functionally, the PRC acts as a critical interface that bridges high-level sensory interpretation, particularly that derived from the ventral visual stream, with the neural machinery responsible for long-term declarative memory formation. It is uniquely positioned to handle complex pattern information and is often considered the principal locus for item-specific Recognition Memory based on familiarity, distinguishing it from the hippocampus, which is traditionally associated with relational or episodic memory based on recollection. This critical integrative role means that the PRC processes the "what" of an experience--the identity of an object or stimulus--before that information is consolidated into enduring memory traces, thereby ensuring that newly encountered objects can be recognized upon future exposure without necessarily recalling the specific context of their initial presentation.

Anatomically, the PRC is part of the parahippocampal region, which collectively includes the PRC, the parahippocampal cortex (PHC), and the entorhinal cortex (EC). This regional organization forms the primary cortical input pathway into the hippocampal formation. Unlike the PHC, which typically processes contextual and spatial information (the "where" of an experience), the PRC receives dense projections conveying highly abstracted, modality-specific information, especially pertaining to the visual characteristics necessary for object identification. The integrity of the PRC is paramount for tasks requiring the discrimination of visually similar items, demonstrating its fundamental role in differentiating distinct perceptual identities within the environment. If this structure is compromised, the ability to mentally separate common objects as unique entities is severely impaired, highlighting its necessary involvement in resolving perceptual ambiguity and encoding object distinctiveness.

The operational definition of the PRC emphasizes its dual functional nature: both perceptual and mnemonic. As a perceptual processing center, it facilitates the resolution of overlapping features among objects, enabling sophisticated Visual Discrimination. As a mnemonic center, it generates the signal of familiarity that allows rapid judgments of prior encounter. These two functions are deeply intertwined; the ability to correctly encode the unique perceptual features of an item is prerequisite for later successful recognition based on familiarity. Consequently, the PRC represents one of the final cortical stages of visual processing before information transitions into the core memory circuit of the MTL, solidifying its status as a gatekeeper for declarative memory formation regarding item identity.

## 2. Etymology and Historical Development

The designation of the perirhinal cortex derives from its anatomical location--it is the cortical area "peri" (around) the rhinal sulcus, a prominent fissure in the temporal lobe of many mammals, including primates. Early neuroanatomical studies utilized cytoarchitectonic mapping, pioneered by figures like Korbinian Brodmann, which characterized the PRC based on the distinct cellular organization (cytoarchitecture) of its layers, primarily identifying it as areas 35 and 36. These classifications established the PRC as a distinct entity from the neighboring entorhinal cortex (Area 28) and the surrounding temporal cortices. Historically, the entire MTL structure was viewed monolithically in relation to memory, largely influenced by the seminal case study of H.M., whose severe anterograde amnesia following bilateral MTL removal underscored the region's absolute necessity for new memory formation.

A pivotal shift in understanding occurred during the 1970s and 1980s, driven by detailed lesion studies in monkeys, notably by researchers such as Mortimer Mishkin and Larry Squire. These studies began to dissect the functional specificity within the MTL components. Critically, these experiments demonstrated that selective lesions of the hippocampus or the amygdala produced different types of deficits than those resulting from lesions encompassing the PRC and the parahippocampal cortex. Specifically, it was the damage to the PRC and the adjacent ventral temporal structures that consistently resulted in profound impairment in delayed non-matching-to-sample tasks, which are standard behavioral measures of visual object recognition memory. This empirical evidence firmly established the PRC's specialized role in recognizing items based on their intrinsic features, separating its function from the hippocampus's spatial and relational processing role.

Further refinement came with the development of the "Dual-Process Theory" of recognition memory in the 1990s and early 2000s. This theory posited that recognition memory is supported by two distinct processes: familiarity (a quick, context-free sense of "knowing") and recollection (a slower, detailed retrieval of contextual specifics). Extensive electrophysiological and neuroimaging evidence accumulated suggesting that the PRC primarily mediates the familiarity component, whereas the hippocampus is central to recollection. This framework solidified the PRC's conceptual role not merely as a relay station, but as an active computational hub where the memory signal of item familiarity is specifically generated. Modern research continues to investigate this distinction, often using functional magnetic resonance imaging (fMRI) and single-unit recordings to confirm the differential activation of the PRC during familiarity-based recognition judgments versus recollection-based retrieval.

## 3. Key Anatomical Characteristics and Inputs

The anatomical location of the PRC makes it an apex structure for sensory processing pathways

converging onto the MTL memory system. It is situated on the ventromedial surface of the temporal lobe, with Area 36 covering the lateral bank of the rhinal sulcus and extending onto the adjacent temporal neocortex, while Area 35 lies medially, lining the sulcus itself. The cellular architecture reveals a relatively simpler structure compared to the six-layered neocortex, particularly in Area 35, which contains fewer layers, reflecting its status as a transitional zone between the neocortex and the archicortex of the hippocampus. This specific anatomical arrangement dictates its functional specialization, allowing it to efficiently consolidate and transfer highly processed sensory information.

A defining characteristic of the PRC is its privileged access to the outputs of the **Ventral Visual Stream**, often termed the "What" pathway. This stream originates in the primary visual cortex (V1) and progresses through increasingly complex processing stages in the occipital and inferior temporal cortices (ITCs). The ITCs, which perform high-level analysis of object features, texture, color, and shape, send massive, direct projections to Area 36 of the PRC. Consequently, the PRC receives visual input that is already highly refined--representing whole objects or complex features rather than simple lines or edges. This concentration of object-related visual information explains why the PRC is so fundamentally involved in visual recognition and object discrimination tasks.

While visual input is dominant, the PRC is truly a polymodal association area, receiving significant afferent projections from multiple sensory and association cortices. It receives input concerning auditory and somatosensory information from the adjacent temporal cortices, as well as crucial modulatory input from the amygdala (related to emotional valence) and the prefrontal cortex (related to executive control and attention). These diverse inputs converge, allowing the PRC to form a comprehensive, multimodal representation of an item, though the visual component remains the most extensively studied and functionally paramount. The integration of these various sensory streams within the PRC is critical for establishing a robust, non-contextual item representation that can be quickly accessed for familiarity judgments.

#### 4. Critical Role in Visual Object Discrimination

The most significant cognitive function attributed to the PRC, particularly within the perceptual domain, is its essential role in resolving **perceptual ambiguity**, a process critical for Visual Discrimination. Visual discrimination involves the ability to correctly identify and distinguish one object from another, especially when the objects share many overlapping features. For instance, distinguishing between two highly similar models of a car, or two subtly different faces, requires fine-grained analysis of specific features and the binding of those features into a unique representation--a computational demand largely met by the PRC.

Research has robustly shown that damage to the PRC leads to a specific type of impairment known as a perceptual memory deficit, where the patient or subject can see the objects clearly, but

struggles immensely when required to distinguish between them, particularly when the distractors are perceptually similar to the target item. The example provided in the source material, illustrating Nancy's inability to discern common objects in her surroundings as separate identities, directly reflects a failure in this discrimination function. This inability is not merely a failure of general memory retrieval; rather, it is a deficit rooted in the foundational encoding process, where the unique item representations necessary to differentiate the objects were never adequately established or maintained by the damaged PRC circuitry.

This discrimination function is often framed within the concept of "pattern separation" applied to item identity. While the dentate gyrus of the hippocampus is known for separating similar spatial or relational patterns, the PRC performs an analogous function for object patterns. It actively computes the differences between incoming sensory data and stored object representations, effectively creating unique neural signatures for items that share many components. This requires a large pool of neurons capable of rapid plastic changes, allowing the brain to adapt item representations based on novel or confusing input. Without this discriminatory capability supplied by the PRC, the perceptual world collapses into an undifferentiated stream of familiar but indistinguishable stimuli, leading to profound recognition failure.

## 5. Functional Connectivity and Memory Pathways

The perirhinal cortex serves as a primary relay node within the declarative memory circuit, mediating the flow of information both into and out of the hippocampal formation. The flow of information is highly organized: high-level sensory input (from the ITCs) first enters the PRC (and PHC), and these areas then project heavily to the Entorhinal Cortex (EC). The EC acts as the final gateway, projecting into the hippocampus proper via the perforant pathway (specifically targeting the dentate gyrus, CA3, and CA1 subfields). This trisynaptic circuit (EC to DG/CA3/CA1) is the classic pathway for episodic memory encoding.

However, the PRC also possesses crucial, direct connections that bypass parts of the full trisynaptic circuit, particularly in relation to familiarity-based memory. Projections from the PRC directly target the deep layers of the EC and bypass the DG entirely, reaching the CA1 subfield of the hippocampus. This shorter, more direct route is hypothesized to be the anatomical substrate for the rapid, non-contextual familiarity signal. When a previously encountered item is presented, the PRC rapidly activates based on the match between the current sensory input and its stored item representation, sending a swift signal of familiarity to the downstream hippocampal areas, thus enabling quick recognition judgments without the need for detailed contextual retrieval.

Furthermore, the PRC maintains extensive reciprocal connections with the prefrontal cortex (PFC), particularly areas involved in working memory and decision-making (e.g., ventrolateral PFC). These connections are essential for the strategic control of memory retrieval and the comparison

process necessary for object discrimination. When a subject attempts to distinguish between two similar objects, the PFC engages the PRC to access and compare the detailed item representations, illustrating that the PRC is not merely a passive storage site but an active component integrated into larger executive and decision-making networks responsible for goal-directed behavior based on recognition.

## 6. The Familiarity Component of Recognition Memory

The PRC is widely recognized as the dominant neural substrate for the "familiarity" component of Recognition Memory. Familiarity is defined as a general, acontextual feeling that an item has been previously encountered--a "knowing" without a specific "remembering." This contrasts sharply with recollection, which involves the retrieval of specific contextual details associated with the original learning event (e.g., where and when the item was seen). Functional neuroimaging and lesion studies consistently demonstrate that while damage to the hippocampus selectively impairs recollection, damage to the PRC results in a disproportionate loss of the ability to make familiarity judgments.

This specialization is thought to arise because the PRC is optimized for representing complex item features, whereas the hippocampus is optimized for representing the spatial and temporal relations \*between\* items and contexts. When an item is encountered, the PRC rapidly integrates its perceptual features and compares the resulting pattern against a memory trace stored locally within its own intrinsic connectivity or surrounding temporal cortices. If a sufficient match occurs, the familiarity signal is generated. This familiarity signal is computationally faster and requires fewer resources than the full reconstruction of an episodic memory, making it highly adaptive for quick decisions about environmental stimuli.

The PRC's ability to sustain representations across time is also crucial for working memory processes that involve object identity. It allows for the maintenance of object representations over short delays, facilitating tasks such as sequential object manipulation or visual matching. This sustained activity in the PRC during delay periods suggests that its function extends beyond mere encoding initiation and contributes actively to maintaining object representations in an accessible state, reinforcing its role as a persistent store for item-specific information necessary for both short-term perceptual discrimination and long-term declarative memory based on familiarity.

## 7. Clinical Implications and Pathophysiology

Disruption of the perirhinal cortex is associated with significant memory and perceptual deficits, primarily manifesting as impairments in visual object recognition and item familiarity. Such damage is frequently observed in clinical conditions affecting the medial temporal lobe. For example, damage caused by temporal lobe epilepsy, viral encephalitis (such as Herpes Simplex

Encephalitis), stroke, or selective neurodegeneration can isolate the PRC and produce highly specific symptoms. These symptoms often include various forms of visual agnosia or profound difficulty performing tasks that require distinguishing between perceptually overlapping stimuli, known as pattern completion failures in the item domain.

In the context of Alzheimer's Disease (AD), the PRC is one of the earliest cortical regions to exhibit pathological changes, specifically the accumulation of neurofibrillary tangles. The progression of AD pathology often begins in the EC and rapidly spreads to the PRC, well before significant involvement of the hippocampus itself. This early pathology in the PRC is thought to underlie some of the initial cognitive complaints in AD patients, particularly difficulties with recognition memory and object identification, demonstrating that the structural integrity of the PRC is intimately linked to the early maintenance of declarative memory function and semantic stability.

The illustrative case mentioned in the source material--where damage to the PRC resulted in Nancy's inability to discern objects as separate identities--exemplifies the classic clinical outcome of PRC lesions. This scenario highlights a breakdown of perceptual memory where objects are registered, but their unique features cannot be sufficiently segregated from one another, leading to a confusion of identities. This deficit is highly specific, often leaving spatial memory (hippocampal function) relatively intact, further supporting the segregation of function within the MTL memory system. Understanding the vulnerability of the PRC is therefore critical for diagnosing and understanding the progression of various neurodegenerative diseases.

## Further Reading

[Perirhinal Cortex \(Wikipedia\)](#)

[Medial Temporal Lobe \(Wikipedia\)](#)

[Recognition Memory \(Wikipedia\)](#)

[Visual Discrimination \(Wikipedia\)](#)