

# Spontaneous Abortion

Authored by  
**mohammad looti**

October 5, 2025

## RECOMMENDED CITATION

mohammad looti (2025). *Spontaneous Abortion*. PSYCHOLOGICAL SCALES. Retrieved from <https://scales.arabpsychology.com/?p=35470>

## Spontaneous Abortion

**Primary Disciplinary Field(s):** Medicine, Obstetrics and Gynecology, Public Health, Reproductive Health

### 1. Core Definition

**Spontaneous abortion**, commonly known as a **miscarriage**, refers to the involuntary termination of a pregnancy before the fetus has reached viability, typically defined as occurring before the 20th week of gestation. This medical event signifies the loss of a fetus or embryo due to natural causes, distinguishing it from an induced abortion, which is the intentional termination of a pregnancy. The threshold of 20 weeks is clinically significant because pregnancies lost after this point are typically termed **stillbirths**, reflecting a different set of etiological factors and clinical management approaches. Spontaneous abortion is a relatively common occurrence, affecting a significant percentage of clinically recognized pregnancies, often before a woman is even aware she is pregnant.

The definition is further refined by gestational age, with early miscarriages (before 12 weeks) being the most prevalent, accounting for approximately 80% of all spontaneous abortions. These early losses are frequently attributed to severe chromosomal abnormalities in the embryo, rendering it unable to develop beyond a certain stage. As pregnancy progresses, the incidence of spontaneous abortion decreases, and the causes become more varied, encompassing a broader range of maternal and fetal factors. Understanding this core definition is fundamental to appreciating the complex biological and psychological aspects associated with pregnancy loss, which extends beyond mere statistical incidence to impact individuals and families profoundly.

### 2. Etymology and Historical Development

The term "abortion" itself derives from the Latin "abortio," meaning "miscarriage" or "untimely birth." Historically, the concept of pregnancy loss has been recognized across cultures, though the medical understanding and terminology have evolved considerably. Ancient texts and medical treatises, from Hippocratic writings to medieval European medicine, described instances of early pregnancy loss, often attributing them to various factors such as maternal health, diet, or even supernatural influences. However, the distinction between spontaneous and induced abortion was not always clearly delineated in lay terms or even early medical discourse, leading to a conflation of natural loss with deliberate termination.

The formal medical distinction and the term "spontaneous abortion" gained prominence with the advancement of modern obstetrics and gynecology in the 19th and 20th centuries. As medical science progressed, particularly with a deeper understanding of human embryology, genetics, and maternal physiology, the mechanisms underlying natural pregnancy loss began to be elucidated.

The ability to confirm pregnancy earlier, coupled with diagnostic tools like ultrasound, allowed for more precise identification and categorization of these events. This historical progression has been crucial in shifting perceptions from often superstitious or moralistic interpretations to a more evidence-based, clinical understanding of spontaneous abortion as a distinct medical condition with identifiable causes and, in some cases, preventable risk factors.

### 3. Pathophysiology and Key Characteristics

The pathophysiology of spontaneous abortion is complex and multifactorial, involving an intricate interplay of genetic, endocrine, anatomical, infectious, and immunological factors. Approximately 50-70% of early spontaneous abortions are attributed to **chromosomal abnormalities** in the embryo or fetus, such as aneuploidy (e.g., trisomy, monosomy) or polyploidy. These errors, often random events occurring during gamete formation or early embryonic cell division, lead to developmental arrest and subsequent expulsion. The body's natural physiological mechanisms recognize these non-viable pregnancies, initiating a process of uterine contractions and cervical dilation to expel the uterine contents.

Beyond chromosomal issues, other key characteristics and underlying mechanisms include structural abnormalities of the uterus (e.g., septate uterus, fibroids), which can impair implantation or fetal growth. Endocrine disorders, such as uncontrolled **diabetes mellitus**, thyroid dysfunction, or **polycystic ovary syndrome (PCOS)**, can create an unfavorable environment for pregnancy maintenance due to hormonal imbalances. Immunological factors, though less understood, are also implicated, where the maternal immune system may mistakenly attack the pregnancy. Additionally, severe systemic infections in the mother can lead to inflammation, fever, and direct damage to the fetus or placenta, triggering pregnancy loss.

Spontaneous abortions are further categorized clinically based on their presentation and progression. A **threatened abortion** involves vaginal bleeding with a closed cervix and a viable pregnancy, with the possibility of progression to complete loss or continuation. An **inevitable abortion** presents with bleeding and cervical dilation, indicating that loss is imminent. An **incomplete abortion** occurs when some, but not all, of the fetal or placental tissue has been expelled from the uterus. A **complete abortion** signifies that all pregnancy tissue has been expelled, and the uterus is empty. A **missed abortion** is characterized by the death of the embryo or fetus, but the tissue remains in the uterus without symptoms of expulsion. Finally, a **septic abortion**, though rare in contemporary medical settings, involves infection of the uterine contents, often complicating an incomplete or missed abortion, posing a severe risk to maternal health.

### 4. Causes and Risk Factors

The causes of spontaneous abortion are diverse and often synergistic, making it challenging to

pinpoint a single culprit in many cases. The most common cause, as previously mentioned, is **chromosomal abnormalities** in the developing embryo or fetus. These genetic errors are typically random and not inherited, reflecting errors during cell division. Advanced maternal age is a well-established risk factor, with the incidence of chromosomal abnormalities and thus miscarriage rising significantly after the age of 35 due to age-related changes in oocyte quality. Paternal age may also play a role, though to a lesser extent.

Maternal health conditions contribute substantially to the risk. Uncontrolled chronic diseases such as **Type 1 or Type 2 diabetes**, severe hypertension, and certain autoimmune disorders (e.g., **systemic lupus erythematosus**, antiphospholipid syndrome) can compromise placental development or fetal well-being. Uterine abnormalities, whether congenital (e.g., bicornuate uterus) or acquired (e.g., large uterine fibroids, severe intrauterine adhesions from previous surgeries), can interfere with implantation or accommodate fetal growth, increasing the risk of loss. Infections, particularly those affecting the reproductive tract or causing systemic illness in the mother (e.g., rubella, toxoplasmosis, cytomegalovirus, HIV), can directly harm the fetus or placenta.

Lifestyle factors and environmental exposures also constitute significant risk factors. The source content correctly identifies **substance abuse**, including alcohol consumption, illicit drug use, and **smoking**, as contributors to miscarriage risk. These substances can have direct toxic effects on the embryo or fetus, impair placental function, or lead to maternal malnutrition. Certain medications, especially those contraindicated in pregnancy, can also be teratogenic or induce uterine contractions, leading to pregnancy loss. Exposure to environmental toxins, such as heavy metals or certain pesticides, has also been implicated, though often requiring high levels of exposure. Obesity, both before and during pregnancy, is another recognized risk factor, potentially due to associated inflammatory states and endocrine dysregulation.

## 5. Clinical Presentation and Diagnosis

The clinical presentation of spontaneous abortion typically involves a combination of symptoms, with **vaginal bleeding** and **abdominal cramping** being the most common and often the first indicators. The bleeding can range from light spotting to heavy flow, sometimes accompanied by the passage of clots or tissue. Abdominal pain is often described as crampy, similar to menstrual cramps, and can vary in intensity and location. Other less specific symptoms may include backache, a decrease in pregnancy symptoms (e.g., nausea, breast tenderness), or the absence of fetal movement if the pregnancy was advanced enough for it to be felt. It is crucial to note that vaginal bleeding in early pregnancy does not always indicate a miscarriage; many women experience spotting and go on to have healthy pregnancies.

Diagnosis relies on a combination of clinical evaluation, laboratory tests, and imaging studies. A physical examination typically includes a speculum examination to assess the amount of bleeding

and determine if the cervix is open or closed, and a bimanual examination to assess uterine size and tenderness. Blood tests are essential to measure **human chorionic gonadotropin (hCG) levels**, the "pregnancy hormone." Serial hCG measurements, taken 48-72 hours apart, can indicate whether the pregnancy is progressing normally, is ectopic, or is failing, as expected hCG levels should double during early pregnancy. A single low or declining hCG level, coupled with symptoms, can suggest a miscarriage.

The definitive diagnostic tool is **ultrasonography**, specifically transvaginal ultrasound in early pregnancy. This imaging technique allows for direct visualization of the gestational sac, yolk sac, embryo, and fetal heart activity. Key ultrasound findings indicative of spontaneous abortion include an empty gestational sac when a fetal pole should be visible (blighted ovum), a fetal pole without cardiac activity, or a gestational sac that is too small for the expected gestational age. In cases of suspected incomplete abortion, ultrasound can also identify retained products of conception within the uterus, guiding management decisions. The combination of symptoms, hCG levels, and ultrasound findings allows clinicians to accurately diagnose the type of spontaneous abortion and plan appropriate care.

## 6. Management and Treatment

The management of spontaneous abortion depends largely on the type of miscarriage, the gestational age, and the patient's preferences and clinical stability. For a **threatened abortion**, management is typically expectant, involving rest, avoidance of strenuous activity, and close monitoring of symptoms and ultrasound findings, as a significant number of these pregnancies will continue successfully. For inevitable, incomplete, or missed abortions, the primary goal is to ensure the complete expulsion of uterine contents and prevent complications such as hemorrhage or infection.

There are three main approaches to managing an inevitable, incomplete, or missed abortion:

**Expectant Management:** This involves waiting for the body to naturally expel the remaining pregnancy tissue. It is suitable for stable patients with no signs of infection and is often preferred by those who wish to avoid medical or surgical intervention. This approach requires patience, as the process can take days to weeks, and close follow-up to ensure complete expulsion.

**Medical Management:** Medications, most commonly **misoprostol** (a prostaglandin analog), are used to induce uterine contractions and facilitate the expulsion of tissue. This method is highly effective, typically occurring within a few hours to days, and can be administered orally or vaginally. It is often chosen for patients who desire a more predictable timeline than expectant management but wish to avoid surgery.

**Surgical Management:** Procedures such as **dilation and curettage (D&C)** are performed to

surgically remove retained products of conception from the uterus. This is a quick and effective method, often preferred for patients with heavy bleeding, signs of infection, or those who prefer a rapid resolution. D&C may also be necessary if expectant or medical management fails.

Irrespective of the management approach, psychological and emotional support is a critical component of care. Experiencing a spontaneous abortion can be a deeply traumatic event, and patients often benefit from counseling, support groups, and reassurance. Follow-up care also includes discussions about future pregnancy planning, as well as addressing any underlying causes that may be identified, such as uterine anomalies or endocrine imbalances. For women with recurrent pregnancy loss (three or more consecutive miscarriages), a thorough diagnostic workup is recommended to identify treatable causes.

## 7. Psychological and Social Impact

The psychological and social impact of spontaneous abortion is profound and often underestimated. For many individuals and couples, a miscarriage represents not just the loss of a pregnancy but the loss of hopes, dreams, and the future they envisioned for their family. The emotional responses can be intense and varied, including **grief**, sadness, anger, guilt, anxiety, and depression. These feelings can be compounded by a sense of isolation, as societal acknowledgment and support for pregnancy loss are often less visible than for other forms of bereavement. The duration and intensity of grief vary widely, influenced by factors such as the gestational age, the number of previous losses, and the individual's coping mechanisms and support system.

Partners also experience significant emotional distress, though their grief may manifest differently and is sometimes overlooked. The shared trauma can strain relationships, while open communication and mutual support are crucial for navigating this difficult period. Socially, there can be pressure to "move on" quickly, which can invalidate the grieving process. Miscarriage can lead to difficulties in subsequent pregnancies, with increased anxiety and fear of recurrence, a phenomenon sometimes referred to as "phantom pregnancy syndrome." Providing adequate psychological support, including access to counseling, peer support groups, and resources, is essential for helping individuals and couples process their loss and promote healing.

## Further Reading

[Miscarriage - Wikipedia](#)

[Stillbirth - Mayo Clinic](#)

[Early Pregnancy Loss \(Miscarriage\) - ACOG](#)

[What is Diabetes? - NIDDK](#)

[Polycystic Ovary Syndrome \(PCOS\) - NICHD](#)

[Septic Abortion - NCBI Bookshelf](#)

[Facts About Birth Defects - CDC](#)

[Type 1 Diabetes - CDC](#)

[What is Lupus? - Lupus Foundation of America](#)

[National Institute on Alcohol Abuse and Alcoholism \(NIAAA\)](#)

[Health Effects of Cigarette Smoking - CDC](#)

[Vaginal Bleeding During Pregnancy - ACOG](#)

[Human Chorionic Gonadotropin \(hCG\) - Mayo Clinic Laboratories](#)

[Fetal Ultrasound - RadiologyInfo.org](#)

[Misoprostol - Drugs.com](#)

[Dilation and Curettage \(D&C\) - ACOG](#)

[Grief and Loss - American Psychiatric Association](#)

ARABPSYCHOLOGY.COM